

Prolactin (PRL) and Its Receptor: Actions, Signal Transduction Pathways and Phenotypes Observed in PRL Receptor Knockout Mice

CHRISTINE BOLE-FEYSOT*, VINCENT GOFFIN*, MARC EDERY, NADINE BINART,
AND PAUL A. KELLY

INSERM Unité 344-Endocrinologie Moléculaire, Faculté de Médecine Necker, 75730 Paris Cedex 15, France

- I. Introduction
- II. PRL
 - A. The PRL/GH/PL family
 - B. PRL gene and primary structure
 - C. PRL tertiary structure
 - D. Extrapituitary PRL
- III. PRL Receptor (PRLR)
 - A. The class 1 cytokine receptor superfamily
 - B. PRLR gene and primary structure
 - C. PRLR tertiary structure
 - D. PRLR binding and activation by PRL
- IV. Distribution of the PRLRs
- V. Biological Functions of PRL
 - A. Water and electrolyte balance
 - B. Growth and development
 - C. Endocrinology and metabolism
 - D. Brain and behavior
 - E. Reproduction
 - F. Immunoregulation and protection
 - G. Actions associated with pathological disease states
- VI. Signal Transduction by the PRLR: Structure-Function Relationships
 - A. The JAK-Stat pathway
 - B. The Ras/Raf/MAP kinase pathway
 - C. Other signaling pathways
- VII. Null Mutation of the PRLR Gene
 - A. Gene cloning, vector construction, and generation of PRLR^{-/-} mice
 - B. PRLR gene expression and PRLR protein in PRLR^{-/-} mice
 - C. Impaired mammary gland development and lactation in heterozygous females
 - D. Heterozygote maternal behavior
 - E. Homozygous females are sterile
 - F. Homozygous male fertility
 - G. Other gene-targeted mutations leading to impaired mammary gland and reproductive function
 - H. Other phenotypes of PRLR^{-/-} mice
- VIII. Summary and Conclusions

I. Introduction

PRL and GH, along with placental lactogens (PLs), form a family of hormones that probably result from the duplication of an ancestral gene. It was early in the 20th century that changes in the histology of the anterior pituitary gland of pregnant women were first noted. French researchers were the first to identify a pituitary factor capable of inducing milk secretion in rabbits (1). American scientists made similar observations, and in addition to naming the new pituitary factor prolactin, showed that PRL was able to stimulate the growth of the pigeon crop sac (2). PRL has now been shown to exist in all vertebrates thus far examined.

Because human GH preparations were lactogenic in conventional bioassays, and because early attempts to separate GH and PRL activities failed, there was some question whether a separate PRL existed in humans. There was strong clinical and histological evidence to suggest that the two hormones were present in humans. Finally, human PRL (hPRL) was successfully isolated and purified (3, 4), which led to numerous subsequent pathophysiological studies.

PRL has more actions than all other pituitary hormones combined. The initial step in the action of PRL, like all other hormones, is the binding to a specific membrane receptor, the PRL receptor (PRLR). Similar to the ligand, the PRLR has also been shown to be a member of the same family as the GH receptor and also part of the larger class of receptors, known as the class 1 cytokine receptor superfamily.

In this review we will briefly discuss the structure of PRL and its family members and the fact that PRL is produced at sites outside the pituitary gland (extrapituitary PRL), and thus may act as a hormone, by the classic endocrine pathway, and as a growth factor, neurotransmitter, or immunoregulator, in an autocrine-paracrine fashion. The structural organization of the PRLR and its complex binding and activation will be described, as well as the tissue distribution of the receptor. The original list of 85 different actions of PRL in vertebrates has been expanded to include more than 300 separate functions of this multifaceted hormone. The signal transduction mechanisms activated after the binding of PRL to the receptor will be described. Finally, the phenotypes associated with the knockout of the PRLR gene in mice will be reviewed. Although this approach does not apply to all reported functions of PRL (seasonal actions, species-specific

Address reprint requests to: Paul A. Kelly, INSERM Unité 344-Endocrinologie Moléculaire, Faculté de Médecine Necker, 156 rue de Vaugirard, 75730 Paris Cedex 15, France. E-mail: kelly@necker.fr

* Equal contributors.

effects, etc.), in many instances the knockout model is useful to identify actions directly associated with PRL or PL and, by comparison with other gene deletions, suggests which actions have been taken over by another hormone or cytokine.

II. PRL

A. The PRL/GH/PL family

Three decades after PRL was identified, the amino acid sequence of sheep PRL (also referred to as lactogenic hormone or luteotropic hormone depending on its biological properties) was determined and shown to be a protein of 199 amino acids (5). At the end of the 1970's, the rapid development of cloning technology allowed the identification of the nucleotide sequence of PRL cDNAs from several species (6). As anticipated from earlier structural studies, the primary structure of PRL appeared closely related to that of two other hormones, GH, also of pituitary origin, and PL, secreted by mammalian placenta (6–8). Today, genetic (7, 8), structural (6, 9), binding (9), and functional (6, 9) studies of these three hormones, as well as the more recently identified somatolactin and PRL-related proteins, have clearly demonstrated that they all belong to a unique family of proteins.

B. PRL gene and primary structure

The gene encoding hPRL is located on chromosome 6 (10). It is composed of five exons and four introns with an overall length of ~10 kb (11). The hPRL cDNA is composed of 914 nucleotides and contains a 681-nucleotide open reading frame encoding a prehormone of 227 amino acids (aa), including a signal peptide of 28 aa (12). The mature hPRL thus contains 199 aa, with a total molecular mass of ~23 kDa.

PRL is present in all vertebrates, and cDNAs encoding PRL from several species have been isolated and sequenced (6, 12–17). With the exception of fish, all PRLs identified so far are 197–199 aa and contain six cysteines forming three intramolecular disulfide bonds (Cys 4–11, 58–174, and 191–199 in hPRL). Fish PRLs are shorter than mammal PRLs and lack a dozen residues at the N terminus, including the first disulfide bridge (16). In tilapia, two distinct PRLs have been isolated, which differ by their length (11 aa), their composition (69% aa identity), and their biological activities (16). These two PRLs presumably result from gene duplication (16) and are likely to reflect a situation unique to fish. Although the primary structure of PRL is highly conserved within a given class [e.g., bovine and human PRLs share 74% aa identity (14), carp and salmon PRLs 77% (18)], PRL sequences from distantly related species show a high degree of divergence [e.g., carp and human PRLs share only 36% of similarity (18)]. Posttranslational modifications of mature PRL, including glycosylation, phosphorylation, or proteolytic cleavage, have been reported and recently reviewed (17, 19).

On the basis of sequence comparisons of tetrapod hormones, Martial and collaborators confirmed the earlier hypothesis formulated by Niall and colleagues (7) that the genes encoding PRL, GH, and PL are evolved from a common ancestral gene (12, 20) and located the divergence of

PRL and GH lineages that occurred some 400 million years ago (12, 13). On the other hand, evolutionary studies including fish hormones suggest that the divergence might have occurred up to 820 million years ago (for discussions, see Ref. 21). Finally, PLs, which are only found in mammals, are believed to have an alternative genetic origin, either the GH lineage (primate PLs) or the PRL lineage (nonprimate PLs) (8, 22).

C. PRL tertiary structure

Secondary structure studies (circular dichroism, etc.) have shown that PRL is an all- α -helix protein and contains almost 50% of α -helices, while the remainder of the protein appears to fold into nonorganized loop structures (23). To date, attempts to determine the three-dimensional (3D) structure of PRL via experimental techniques (x-ray, nuclear magnetic resonance) have been unsuccessful. However, taking advantage of the structure/function similarities between PRL and GH (see above), we have recently determined the 3D structure of hPRL using the homology modeling approach (24) based on the crystallographic coordinates of porcine (p) GH (25). As anticipated, hPRL is predicted to fold in a four-helix bundle and to share with GHs the particular up-up-down-down connectivity of the α -helices (9, 24–26) (Fig. 1A).

D. Extrapituitary PRL

In addition to being synthesized and secreted by lactotrophic cells of the anterior pituitary gland, PRL is also produced by numerous other cells and tissues. The subject of extrapituitary PRL has recently been reviewed (27) and thus will not be described here in detail.

In addition to the anterior pituitary gland, PRL gene expression has been confirmed in various regions of the brain, decidua, myometrium, lacrimal gland, thymus, spleen, circulating lymphocytes, and lymphoid cells of bone marrow, mammary epithelial cells and tumors, skin fibroblasts, and sweat glands (reviewed in Ref. 27). PRL can thus be found in several fluid compartments in addition to serum, such as cerebrospinal fluid, amniotic fluid, tears, milk, follicular fluid, and sweat. Interestingly, hypophysectomized rats retain ~20% of biologically active PRL in the circulation, which increases to ~50% of normal levels with time. Neutralization of circulating PRL with anti-PRL antibodies results in immune dysfunction and death (28), suggesting that extrapituitary PRL is important and, under some circumstances, can compensate for pituitary PRL.

Pituitary PRL acts via a classic endocrine pathway, *i.e.*, it is secreted by a gland, transported by the circulatory system, and acts on target cells at some peripheral sites via specific receptors located on the plasma membrane. The PRL that is produced by many different cell types can act in a more direct fashion, *i.e.*, as a growth factor, neurotransmitter, or immunomodulator, in an autocrine or paracrine manner. Thus, locally produced PRL can act on adjacent cells (paracrine) or on the PRL-secreting cell itself (autocrine). Using paracrine or autocrine mechanisms, it would thus be possible to activate many of the actions associated with PRL without ever affecting the circulating concentration of the hormone.




FIG. 1. A, Ribbon representation of the predicted 3D structure of hPRL, modeled on the basis of the crystallographic structure of porcine GH (24, 25). hPRL is predicted to adopt the four-helix bundle folding described for GHs (24, 26, 77). Location of binding sites 1 and 2 (see text) is indicated. Side chains of amino acids involved in binding site 1, as deduced from mutational studies (9), are represented. B, Ribbon representation of the 3D x-ray structure of a monomer of the human PRLR ECD (62). The ECD folds in a β -sandwich formed by two antiparallel β -sheets (see text). N- and C-terminal ends are indicated by N and C, respectively. This figure was kindly provided by Drs. P. Elkins and A. M. de Vos. Note that the structures depicted in panels A and B are not at the same scale (see Ref. 62). C, PRLR activation by PRL-induced dimerization. Hormone binding to PRLR is sequential. First, the hormone (H) interacts with the receptor (R) through its binding site 1 (see Fig. 1A), forming an inactive H_1R_1 complex. Then, the hormone binds to a second receptor through its site 2, which leads to receptor homodimerization and formation of an active H_2R_2 complex. Hormone analogs whose binding site 2 is sterically blocked are unable to induce receptor homodimerization and are thus inactive; since they still bind to the receptor through site 1, they behave as antagonists of wild-type hormones (9).

III. PRL Receptor (PRLR)

A. The class 1 cytokine receptor superfamily



More than two decades ago, the PRLR was identified as a specific, high-affinity, membrane-anchored protein (29–32). In 1988, the cDNA encoding the rat PRLR was isolated in our laboratory (33) and, as is true for their respective ligands, receptors for PRL and GH (GHR) are also closely related (33–35). Both are single-pass transmembrane chains and, despite a relatively low degree (~30%) of sequence identity, they share several structural and functional features (35–38). In the early 1990's, sequence comparison with newly identified membrane receptors led to the identification of a new family of receptors including both PRLR and GHR (35, 39, 40). Termed class 1 cytokine receptors, this superfamily in-

cludes receptors for several interleukins, granulocyte-colony stimulating factor (G-CSF), granulocyte macrophage-colony stimulating factor (GM-CSF), leukemia inhibitory factor (LIF), Oncostatin M (OM), erythropoietin (EPO), thrombopoietin (TPO), gp130, and the obesity factor leptin (41–45). Although all these membrane chains are apparently genetically unrelated, they contain stretches of highly conserved amino acids, both in the extracellular and the intracellular domains. These conserved features are described below with respect to the PRLR.

B. PRLR gene and primary structure

The gene encoding human PRLR is located on chromosome 5 (p13→14) and contains at least 10 exons for an overall length exceeding 100 kb (46). Contrary to PRL, for which a single transcript encodes a unique mature protein, multiple isoforms of membrane-bound PRLR resulting from alternative splicing of the primary transcript have been identified in several species (33, 47–51). These different PRLR isoforms differ in the length and composition of their cytoplasmic tail and are referred to as short, intermediate, or long PRLR with respect to their size (Fig. 2A) (For review, see Refs. 35, 37, and 38). For example, in rat, the PRLR isoforms contain 291 (short), 393 (intermediate), or 591 (long) aa. In mice, one long and three short isoforms have been identified, the short forms only differing by a few amino acids in the C-terminal part of their cytoplasmic tail (51, 52). In addition to the membrane-anchored PRLR, soluble isoforms have also been identified (PRL binding protein, or PRLbp), but whether they result from alternative splicing of the primary mRNA or proteolytic cleavage of membrane-bound PRLR (or both) is uncertain (53–55). In all cases, however, the extracellular, ligand-binding domain is identical, whatever the isoform. Detailed description of the various PRLR isoforms from different species has been provided in previous reviews (35, 37, 38, 56), and this aspect is thus not developed in this review, except when required.

1. *The extracellular domain (ECD)*. Most of the sequence similarities between cytokine receptors are found within their ECD. Typically, a cytokine ECD is composed of a domain of ~200 aa, referred to as the cytokine receptor homology (CRH) region (45). The CRH can be divided into two subdomains of ~100 aa (referred to as D1 and D2), each showing analogies with the fibronectin type III module (35, 40, 45). Although some cytokine receptors contain additional domains, it seems that ligand interactions are primarily driven by the conserved fibronectin-like domains (45, 57). In rat and human, the PRLR ECD encompasses the 210 amino-terminal residues, whatever the isoform considered (33, 35, 48). Two highly conserved features are found in the cytokine receptor ECDs: the first is two pairs of disulfide-linked cysteines in the N-terminal subdomain D1 (Cys12–Cys22 and Cys51–Cys62 in hPRLR), and the second is a pentapeptide termed "WS motif" (Trp-Ser-any amino acid-Trp-Ser) found in the membrane-proximal region of the C-terminal subdomain D2 (Fig. 2A). The functional importance of these features is discussed below (Section III.D). The case of avian PRLRs is atypical since, at least in pigeon and chicken, the PRLR ECD is du-

A**B**

plicated and contains two highly homologous CRH regions (58). The additional N-terminal module does not seem to play any functional role since its deletion has no significant effect on the ligand-binding affinity, ligand specificity (58), or signal transduction (59) of the pigeon PRLR.

2. The transmembrane domain. Like all cytokine receptors, the PRLR is a single-pass transmembrane chain. The transmembrane domain is 24 aa long (aa 211–234 in rat PRLR). The involvement of this region (or of any crucial amino acid within this domain) in the functional activity of the receptor is unknown.

3. The intracellular domain. The cytoplasmic domain of cytokine receptors displays more restricted sequence similarity than the ECD. Two regions, called box 1 and box 2 (35, 60), are relatively conserved. Box 1 is a membrane-proximal region composed of 8 aa highly enriched in prolines and hydrophobic residues (aa 243–250 in PRLR; Fig. 2A). Due to the particular structural properties of proline residues, the conserved P-x-P (x = any amino acid) motif within box 1 is assumed to adopt the consensus folding specifically recognized by transducing molecules (see below). The second consensus region, box 2, is much less conserved than box 1 and consists in the succession of hydrophobic, negatively charged, then positively charged residues (aa 288–298). While box 1 is conserved in all membrane PRLR isoforms, box 2 is not found in short isoforms (35, 38).

In a recent study (61), we have identified, within the cytoplasmic domain of the short PRLR, two motifs required for receptor internalization. The first involves a dileucine motif (aa 259–260); the second contains a tetrapeptide predicted to fold in a β -turn (aa 276–279). Interestingly, the long PRLR isoform, which is less efficiently internalized than the short form, lacks the putative β -turn motif (61).

C. PRLR tertiary structure

The 3D structure of genetically engineered hPRLR ECD (*i.e.*, hPRLbp) has been determined by crystallographic analysis (62) (Fig. 1B). Each fibronectin-like subdomain (D1 and D2) contains seven β -strands that fold in a sandwich formed by two antiparallel β -sheets, one composed of three strands referred to as strands A, B, and E, and the other composed of the four remaining strands termed C, C', F, and G (26, 38, 45, 57, 62). Both subdomains are linked by a small four-residue polypeptide (26, 62). As anticipated from sequence

comparison (40), this folding pattern is likely to be shared by several, if not all, cytokine receptors, since it has also been described for the ECDs of the hGH receptor (26) and the EPO receptor (63, 64) as well as for the α -chain of the interferon (IFN)- γ receptor, a class 2 cytokine receptor (65). To the best of our knowledge, no structural data have been reported yet for the cytoplasmic domain of any cytokine receptor, including the PRLR.

D. PRLR binding and activation by PRL

No exhaustive information on the amino acids of the PRLR ECD interacting with PRL is yet available. Actually, two mutational studies performed in our laboratory focused on some features conserved in cytokine receptor ECDs, including the two pairs of disulfide-bonded cysteines and the WS motif (see above, *Section III.B*) (66, 67). In agreement with similar studies performed on the GHR (68), mutation of any of these conserved cysteines leads to impaired structural and functional properties of the receptors (66), although only amino acids bordering the first, but not the second, disulfide bridge are likely involved in ligand binding (67). Despite the fact that structural data clearly indicate that the WS motif in both GHR (26) and PRLR (62) is located away from the ligand-binding interface, mutations within this conserved feature are detrimental to binding affinity (67, 69) (for review, see Ref. 38). Actually, functional studies of several cytokine receptors (69–72) have suggested that the WS motif is probably required for correct folding and cellular trafficking rather than for ligand binding itself (for discussions, see Ref. 38). Finally, in addition to these features typically conserved in cytokine receptors, we have also suggested that two tryptophans (Trp72 and Trp139) of the PRLR are involved in PRL binding. This hypothesis is consistent with the 3D structure of the two homologous complexes, hGH-hGHbp and hGH-hPRLbp (26, 62), and suggests that these two tryptophans represent a specific feature of the ligand-receptor interactions within the PRL/GH family (24, 38). The three asparagine-linked glycosylation sites present in the ECD of the PRLR do not appear to be involved in ligand binding (62, 73, 73a).

Although stoichiometric analysis of the interaction between different PRLR ECDs and lactogenic hormones achieved 1:1 (74) or 1:2 (75) complexes depending on the species involved, dimerization of the PRLR upon ligand binding has now been clearly established after different approaches. First, we have shown that at least two regions of

FIG. 2. A, Schematic representation of soluble (human) and membrane (rat) isoforms of the PRLR (33, 47, 50, 54). Although the mechanism of PRLbp generation remains unclear (alternative splicing or proteolysis or both), an mRNA encoding a soluble PRLbp of 206 aa has been isolated in the human breast cancer cell line BT-474 (54). All forms have identical extracellular, ligand-binding domains. Subdomain D1 contains two pairs of disulfide bonded-cysteines (C-C) and subdomain D2 contains the WS motif (*green box*), two characteristic features of the cytokine receptor superfamily. Box 1 (*orange box*) is found in the cytoplasmic domain of all membrane isoforms. In rat, the intermediate PRLR (only found in Nb2 cells) differs from the long isoform by a 198-aa deletion in the cytoplasmic domain (aa 323–520). Otherwise, the short PRLR is identical to both other isoforms up to residue 261, after which its sequence differs (*light blue box*). Cytoplasmic tyrosine residues are indicated. B, Structure-function relationships of the long PRLR cytoplasmic domain. Box 1 is required for JAK2 binding; whether this interaction is direct or mediated by an adapter is unknown. The di-leucine motif (aa 259–260), identified in the short PRLR, is presumably involved in internalization of all PRLR isoforms. Six tyrosines (of the nine present in rat PRLR) are potentially phosphorylated. The most C-terminal (Y580), required for Stat5 activation, is proposed to be the major binding site of this Stat protein. Y479 and Y473 can also activate Stat5, although to a lesser extent; these may be Stat5-binding sites of lower affinity. Stats 1 and 3 are likely to interact with membrane-proximal regions of the receptor complex; candidates are Y309 on the receptor or tyrosines to be defined within JAK2. The membrane-proximal region that is common to all PRLR isoforms is required for interaction with and/or activation of JAK2, Fyn, and MAP kinases as well as for activation of cell proliferation and transcription of milk protein genes.

hPRL are involved in the binding of the hormone to the PRLR. The first, referred to as binding site 1, encompasses several residues belonging to helices 1 and 4 (24, 76–78), while the second, termed binding site 2, involves helices 1 and 3 (24, 79, 80) (Fig. 1A). Detailed analysis of individual residues required for tight receptor binding has been reported in a recent issue of *Endocrine Reviews* (9). Second, analysis of PRLR ECD-lactogenic hormone complexes using surface plasmon resonance technology (BIAcore, Pharmacia & Upjohn AB, Stockholm, Sweden) has demonstrated the formation of 1:2 complexes (81). It is likely that the very rapid dissociation of 1:2 complexes for 1:1 complexes has prevented their identification by classic gel filtration experiments (74), as well as by the two-hybrid approach (82). Third, elucidation of the events occurring upon PRL-induced activation of membrane-bound PRLR has resulted from the close analysis of the shape of experimental curves obtained with point-mutated hPRL analogs in PRL-responsive bioassays performed over a wide range of hormone concentrations (reviewed in Ref. 9). As first described for the closely related GHR (45, 83, 84), activation of the PRLR involves ligand-induced sequential receptor dimerization (Fig. 1C). In a first step, interaction of PRL binding site 1 with one PRLR occurs and leads to the formation of an inactive $H_1:R_1$ (one hormone, one receptor) complex. Formation of this complex appears to be a prerequisite for PRL binding site 2 to interact with another PRLR, thereby achieving an active trimeric complex ($H_1:R_2$), composed of one hormone and one receptor homodimer. In agreement with this model, hPRL analogs carrying a disruptive mutation in binding site 2 are inactive (since they are unable to induce PRLR dimerization) but display antagonistic properties due to their ability to block the receptor in the inactive $H_1:R_1$ stoichiometry (9, 37, 80).

IV. Distribution of the PRLRs

PRL-binding sites or receptors have been identified in a number of cells and tissues of adult mammals. The expression of short and long forms of receptor have been shown to vary as a function of the stage of the estrous cycle, pregnancy, and lactation (85–88). As can be seen in Table 1, PRL-binding sites or receptors are widely distributed throughout vertebrates. There was, however, very little information on the expression of this receptor during fetal development. To that end, we have recently determined the cellular distribution and developmental expression of the PRLR in the late gestational fetal rat by *in situ* hybridization, immunocytochemistry, and radioligand binding (89). Sense and antisense strand probes were prepared encoding the long and short isoforms of the rat PRLR and hybridized to various fetal tissues obtained at the end of pregnancy (days 17.5 to 20.5). These studies showed that the mRNA encoding the short and long isoforms was widely expressed in tissues from all three germ layers: in addition to the classic target organs of PRL, tissues not known previously to contain PRLRs, such as olfactory neuronal epithelium and bulb, trigeminal and dorsal root ganglia, cochlear duct, brown adipose tissue, submandibular glands, whisker follicles, tooth primordia, and proliferative and maturing chondrocytes of

TABLE 1. Distribution of PRL binding sites in vertebrates

Cell or tissue	Cell or tissue
Central nervous system	Kidney
Brain	Cortex
Cortex	Bladder (fish, reptiles, amphibians)
Hippocampus	Lymphoid tissue
Choroid plexus	Spleen
Striatum	Thymus
Cochlear duct	Nurse cells
Corpus callosum	Epithelial cells
Hypothalamus	Lymphocytes
Astrocytes	T
Glial cells	B
Retina	Macrophages
Olfactory system	Ganglia
Ganglia	Intestinal cells
Pituitary	Reproductive system
Anterior lobe	Female
Intermediate lobe	Ovary
Adrenal cortex	Ova
Skin	Granulosa cells
Epidermis	Thecal cells
Hair follicle	Corpus luteum (luteal cells)
Sweat gland	Oviduct
Bone tissue	Mammary gland
Chondrocytes	Epithelial cells
Cartilage	Milk*
Osteoblasts	Tumors
Gills (fish and larval amphibians)	Crop sac (birds)
Lung	Uterus (endometrium)
Heart	Placenta
Cardiac muscle	Amnion
Atria	Male
Skeletal muscle	Testis
Adipocytes (birds)	Germ cells
Brown adipose tissue	Spermatozoa
Liver	Leydig cells
Hepatocytes	Sertoli cells
Kupffer cells	Epididymis
Submandibular gland	Seminal vesicle
Submaxillary gland	Prostate
Pancreas	
Islet of Langerhans	
Gastrointestinal tract	
Esophagus	
Stomach	
Intestine	
Duodenum	
Jejunum	
Ileum	
Colon	

* Membrane and soluble forms.

developing bone, also expressed PRLR. There was also a high level of expression of receptor mRNA in the fetal adrenal cortex, gastrointestinal and bronchial mucosae, renal tubular epithelia, choroid plexus, thymus, liver, pancreas, and epidermis.

To complement the *in situ* studies, immunohistochemical studies using monoclonal anti-PRLR antibodies clearly demonstrated the distribution of PRLR immunoreactivity was similar to that of the mRNA, strongly suggesting that the receptor protein is expressed in the developing fetus. The functional activity of the PRLRs was established by the demonstration of specific rat PL II binding sites in fetal adrenal cortex, renal tubules, small intestinal villi, pancreatic ductules and islets, hepatic parenchyma cells, choroid plexus

ependymal cells, fetal lung, and thymus. The level of PRLR mRNA and protein actually increased between days 17.5 and 20.5 of pregnancy in a number of tissues, including the adrenal, pancreas, small intestine, pituitary, thymus, liver, and submandibular gland. These results suggest that lactogenic hormones such as PRL and PLs may play important roles in fetal and neonatal development (89).

Since the PRLR is expressed at relatively low levels in the olfactory bulb of the adult rat, but is easily detected in late pregnancy in the fetal rat, we decided to investigate the ontogenesis of PRLR expression in the olfactory system, again using *in situ* hybridization and immunohistochemistry (90). At embryonic day 12.5 (e12.5), mRNAs encoding the long and short isoforms of the PRLR were detected in the medial and lateral nasal processes, the epithelial lining of the olfactory pit, and the neuroepithelium lining of the cerebral ventricles, in the region of the rhinencephalon. PRLR mRNA was also highly expressed in the frontonasal mesenchyme and the mesenchymal tissue underlying the developing brain and in the interpeduncular fossa. Once again, the PRLR immunoreactivity was similar to that of mRNA, suggesting that the PRLR gene was translated in lactogen binding sites or receptors in the developing embryo. As pregnancy advanced, the receptor was expressed intensely, albeit discontinuously, in the olfactory system. Receptor expression was also seen in the cartilage primordia of the ethmoid, sphenoid, temporal, and mandibular bones. Although the PRLR was expressed in the vomeronasal organ, it was limited to the luminal epithelial surface.

It was not until embryonic day 18 that PRLR mRNA and protein was detected in the olfactory bulb. The highest level of expression was seen in the periventricular neuroepithelium. Thereafter, strong staining was observed in the mitral and tufted cell neurons and the sensory neuronal cell bodies of the olfactory epithelium. This high level of expression continued until neonatal day 5. Interestingly, PRLR expression was also found in the mitral cells of the olfactory bulb of the lactating rat, although the levels appear to be much lower than those seen in the fetal and neonatal rat. These studies suggest novel roles for lactogenic hormones in olfactory differentiation and development and may provide new mechanisms by which lactogenic hormones may regulate neonatal behavior and maternal-infant interactions (90).

V. Biological Functions of PRL

PRL was originally isolated by its ability to stimulate mammary development and lactation in rabbits and soon thereafter to stimulate the production of crop milk in pigeons (1, 2). PRL was shown also to be luteotrophic, that is to promote the formation and action of the corpus luteum (91). Subsequently, a number of additional activities have been associated with this hormone in various vertebrate species. In the now classic reviews by Nicoll and Bern (92) and Nicoll (93), 85 different biological functions of PRL were subdivided into five broad categories: 1) reproduction, 2) osmoregulation, 3) growth, 4) integument, and 5) synergism with steroids. Although these reviews were not exhaustive, the authors tried to consider only relevant effects and disregard those of un-

certain validity. This elevated number of biological actions associated with PRL exceeded by far all of the reported actions of the other anterior pituitary hormones combined.

Since the publication of these reviews in 1972 and 1974, numerous other biological functions of PRL have been identified. This section attempts to deal with the now classic functions of PRL and incorporate the more recent findings in the compilation of the actions of this multifaceted hormone. In addition, we decided to modify the categories originally reported, since there was some overlap, and more importantly, the section on reproduction was, in our estimation, too vast. We have thus divided the actions of PRL into the following categories: 1) water and electrolyte balance, 2) growth and development, 3) endocrinology and metabolism, 4) brain and behavior, 5) reproduction, and 6) immunoregulation and protection. We have described actions and cited references dealing with especially well known actions of PRL in lower species, even though these actions may not be seen in mammals. Such actions of PRL may have been lost with evolution or may only be seen in higher animals during certain stages of development. In addition, certain actions of pituitary-derived PRL (endocrine) may be taken over by locally produced PRL (autocrine or paracrine, see Section II.D).

A. Water and electrolyte balance

Regulation of salt and water balance is an essential aspect of homeostasis for most organisms. This is especially true for animals living in environments that desiccate them (land or seawater) or that inundate them with water or leach out salts (fresh water). PRL is clearly involved in water and electrolyte balance in almost all classes of vertebrates, although these effects are more difficult to demonstrate in mammals, in spite of the fact that PRLRs are present in kidney, as well as other tissues involved in salt balance.

Osmoregulatory problems are greatest for fish that migrate between fresh and seawater. In a marine environment, fish drink large amounts of water to replace water lost osmotically through the gills. They absorb ions from seawater and excrete them by active transport through the gills. Thus, marine fish can absorb water from the gut and replace what is lost. Fish excrete a minimal volume of urine (93).

In fresh water, the body fluids are hypertonic to the external environment and thus must cope with the problem of water inundation and the loss of ions that diffuse out through the gills. Adaptation to the freshwater habitat necessitates a reduction of gill permeability to salt loss and changes the transport process from active excretion to active uptake. The kidney eliminates the increased water by increasing the rate of urine flow (93).

As depicted in Table 2, PRL plays a major role in regulating water and electrolyte balance through the gill and kidney of many fish and has been referred to as a fresh water-adapting hormone. PRL clearly reduces Na^+ loss or efflux (94) and water uptake or permeability (95) in the gill and increases extracellular volume (104) in the kidney. Na^+ reabsorption is also enhanced in the bladder of fish and amphibians (106, 107). Electrolyte changes and water drive are other effects of PRL in amphibians (109, 110). Birds eliminate excess salt by increasing secretion of the nasal salt gland, which is also

TABLE 2. Water and electrolyte balance

Organ or Target	Effect	Animal class	Reference
Gills	↓ Na ⁺ loss (efflux)	Fish	(94)
	↓ Water uptake (permeability)		(95)
	↑ Mucus cell size and number		(96)
	Inhibition of Na ⁺ -K ⁺ -ATPase		(97)
	Stimulation of Ca ²⁺ -ATPase		(98)
Kidney	↓ Ca ²⁺ loss (efflux)	Fish	(99)
	↑ Size of renal glomeruli		(100)
	Hypertrophy of renal tubular cells		(101)
	Stimulation of Na ⁺ -K ⁺ -ATPase	Fish, mammals	(97, 102)
	↓ Na ⁺ and K ⁺ excretion	Mammals	(103)
	↑ Extracellular volume	Amphibians	(104)
	↑ Plasma urate	Birds	(105)
	↑ Urine flow rate		(105)
	↑ Na ⁺ reabsorption	Fish, amphibians	(106, 107)
	↑ Mucus cell size and number	Fish	(108)
Bladder	Electrolyte change and water drive	Amphibians	(109, 110)
Skin	↓ Na ⁺ and Cl ⁻ in sweat	Mammals	(111)
Sweat gland	↑ Water and salt absorption:	Mammals	(112)
Intestine	Duodenum: Na ⁺ , Ca ²⁺		(113)
	Jejunum: Na ⁺ , K ⁺ , Ca ²⁺		(114)
	Ileum: Na ⁺ , Cl ⁻ , K ⁺ , Ca ²⁺		(115)
	Colon: Na ⁺ , Cl ⁻		(116)
	↑ Secretion (elimination of excess salt)	Birds	(117)
Nasal salt gland	↑ Ca ²⁺ , Na ⁺ , K ⁺ , Cl ⁻ in flushings	Mammals	(118)
Uterus	↓ Fluid volume in amnion	Mammals	(119)
Placenta			
Pathology			
Placenta	Polyhydramnios	Mammals	(120)
Sweat gland	Cystic fibrosis	Mammals	(121)

stimulated by PRL (117). Although the effects on salt and water balance are much less clear than in fish and amphibians, in mammals PRL has been shown to reduce renal Na⁺ and K⁺ excretion (103) and to stimulate Na⁺-K⁺ adenosine triphosphatase (ATPase) (102). In addition, PRL decreases Na⁺ and Cl⁻ in sweat (111) and increases water and salt absorption in all regions of the intestine (112). Finally, PRL induces a reduction in fluid volume in the amnion (119).

B. Growth and development

A large number of the reported effects of PRL are associated with growth and development. Many of these are seen in lower vertebrates, but more recent data confirm that cellular proliferation is also one of the important functions of PRL in mammals. The varied functions associated with the category are summarized in Table 3.

1. Body growth. Although PRL and GH are produced by cells of the anterior pituitary that have a common stem cell, there are clear and distinct functions of these two hormones. Thus, in humans that lack GH or do not have a functional GHR, dwarfism is always observed, in spite of the fact that the pituitary produces a functionally active PRL. There have been isolated reports of a somatogenic effect of PRL on body weight in male rats (123) and a small effect on postnatal body growth (124), but most investigators fail to see any consistent effect on overall body size. Finally, in mice lacking the PRLR gene, and thus any functional PRLR (see Section VII), there is no modification in overall body length (214), which argues strongly against a direct effect of PRL on body growth.

Interestingly, Ames dwarf mice lacking GH, PRL, and TSH have a longer lifespan than normal mice, in spite of the fact

that these animals show some characteristics of reduced immune function (215). Whether this extended life is due to deficiency of the GH/insulin-like growth factor-I (IGF-I) pathway or to PRL deficiency remains to be investigated. The PRLR knockout mouse (see Section VII) represents a good model by which to determine whether the PRL pathway is directly involved in aging (214).

There appears to be extensive overlap in many of the biological functions of PRLs and GHs in many lower vertebrates. In birds, for example, an increase in body weight has been observed in PRL-treated males that was significantly higher than that of control animals (122).

2. Growth and metamorphosis. In amphibians, PRL is best known for its antimetamorphic effects. In larvae, PRL has been shown to increase the growth of gills and caudal fin (127, 128) and to increase tail length (126, 129) and is thus considered to be a larval GH. At the premetamorphic stages, PRL reduces growth of the hind legs (130, 131) and prevents resorption of the tail (132–134), both effects being mediated by a reduction in thyroid hormone receptor autoinduction. Some species of fish also undergo metamorphosis. For example, in the Japanese flounder, injections of ovine PRL antagonized the stimulatory effect of T₃ on the resorption of the dorsal fin rays, whereas ovine GH had no effect (125). Finally, in amphibians, PRL induces a metamorphosis of sodium channels (139, 140) and, in mammals, of visual pigments in the retina (200).

3. Cell proliferation. As seen in Table 3, many of the actions of PRL are associated with an effect on cell proliferation and development. One of the major targets is skin. In reptiles and amphibians, PRL promotes molting of the epidermis (135).

PRL stimulates skin melanocyte growth in fishes and mammals (142, 143), and keratinocyte growth in mammals (144). In birds, PRL induces defeathering and epidermal growth of the incubation patch (136, 141). Finally, in deer and goats, PRL is responsible for seasonal changes in pelage growth cycles (145, 146).

Proliferation of the epithelial cells of the crop sac in pigeons was one of the early functions associated with PRL, and crop sac growth still remains the official biological assay of this hormone (147, 148).

Although the liver is not an organ that undergoes rapid proliferation, there have been some reports suggesting that PRL plays an important role in the turnover of hepatocytes. Initial studies showed that there was a striking increase in the number of mitotic figures in livers of transgenic mice expressing hGH, which binds equally well to PRL and GH receptors. To determine whether the observed effect was due to the somatogenic or lactogenic activity of hGH, hepatocytes were isolated from 20-day-old rats and cultured in the presence of rat GH or rat PRL. Interestingly, these cells from young animals, in contrast to normal hepatocytes from adult animals, proliferate in culture. Rat PRL was able to significantly increase the number of mitotic figures in these cells, in comparison to rat GH, which actually slightly decreased proliferation (154). This suggests a functional role for PRL in hepatocyte growth, at some time during development.

Although for many years no clear function could be associated with the presence of the relatively high expression of PRLRs in liver, examination of Table 3 reveals that there are now a number of factors activated by PRL in liver: these include protein kinase C (PKC), diacylglycerol, mitogen-activated protein (MAP) kinase, and phosphoinositide turnover (155–158). In addition, several growth-related genes are induced in liver by PRL, such as IGF-I, ornithine decarboxylase (ODC), *c-myc*, *c-fos*, *c-jun*, and *c-src* (153, 156, 159, 162). Although IGF-I is normally associated with GH stimulation, there may be some instances in which the expression of this growth factor is more responsive to PRL. The ODC gene is also induced and the protein activated in a number of other tissues, including heart, kidney, muscle, adrenal, gonads, and prostate (153, 178, 192, 195).

The proliferation of many cells not normally associated with a direct effect of PRL has also been reported. This hormone has been shown to induce an increase in the size of the intestinal mucosa (171, 172) and proliferation of vascular smooth muscle (176, 177), of β -cells of the pancreas (180–182), of pituitary GH3 cells (187), of human benign prostate hypertrophy epithelial cells (193), of astrocytes (198), and of various cells of the immune system (202, 203).

4. Differentiation and development. PRL has also been suggested to have some functional activity in various developmental processes. In addition to its antimetamorphic properties, PRL induces maturation of the lung and surfactant production (149, 152), differentiation of preadipocytes (179), maturation of germ cells (188, 189), and tuberoinfundibular hypothalamic dopamine development (199).

5. Tumor growth. Finally, PRL has been associated with certain forms of tumors and may be directly or indirectly in-

involved in tumor growth. These observations are summarized in Section V.G on actions associated with pathological disease states. Interestingly, the Moloney murine leukemia virus integration-2 locus colocalizes to the same region of chromosome 2 (rat) and 15 (mouse) as the PRLR and GHR. In one rat T cell lymphoma line, the PRLR gene was activated by provirus integration into the PRLR promoter, rendering the lymphoma sensitive to PRL (210).

C. Endocrinology and metabolism

The endocrine and metabolic effects of PRL not related to reproduction are summarized in Table 4. PRL has been shown to effect energy metabolism by modulating ATPase activity in monkey brain: Na^+ - K^+ -dependent ATPase was stimulated while Mg^{2+} and Ca^{2+} -dependent ATPases were reduced in neural as well as glial cells (216).

PRL has marked effects on lipid metabolism. In birds, it augments lipoprotein lipase activity in adipocytes (221), although this effect is not seen in mammals, as the adipocyte does not have PRLRs. In mammals, PRL stimulates phospholipid synthesis in the fetal lung (149) and lipoprotein lipase activity in liver (217). Endogenous and exogenous organic compounds, such as cholesterol, bile salts, drugs, and metabolites that cannot be handled by the kidney, are secreted into bile and eliminated. The active transport of bile acids from the plasma to the bile canaliculus, followed by passive movement of water, is an important determinant of bile flow and one of the major hepatic functions. The transport of bile acids such as taurocholate across the basolateral plasma membrane involves a Na^+ - K^+ ATPase. PRL has been shown to increase bile secretion (218) and the mRNA encoding the Na^+ -taurocholate cotransport polypeptide and hepatic Na^+ -taurocholate cotransport (219, 220).

PRL has been reported to affect carbohydrate metabolism in several vertebrate classes, including hyperglycemic/diabetogenic actions. PRL has a differential effect on the activities of enzymes involved in the Embden-Meyerhoff pathway and the hexose monophosphate shunt in neural and glial cells of male monkeys (222). In addition, PRL at physiological concentrations produced a 4-fold increase in glycogen phosphorylase- α activation in isolated hepatocytes (223). Finally, PRL is known to have direct effects on pancreatic function, increasing insulin secretion (185, 224, 225), decreasing glucose threshold for insulin secretion (226), and increasing glucokinase and glucose transporter 2 (227).

A direct action of PRL on adrenal steroidogenesis has been reported. Specifically, PRL is reported to increase adrenal androgens, dihydroepiandrosterone and dihydroepiandrosterone sulfate (229) as well as cortisol and aldosterone (228). Interestingly, a stimulatory effect on 21-hydroxylase activity has also been reported (230). Although there are a number of reports in the literature suggesting that PRL is able to stimulate adrenal catecholamine synthesis directly (234, 235), since there are no receptors in adrenal medullary cells (see Table 1), the effect may be mediated via an increased secretion of adrenocortical hormones stimulated by PRL. In skin, PRL has been shown to increase the expression of type IV 3 β -hydroxysteroid dehydrogenase (231).

Although PRL signal transduction is covered in another

TABLE 3. Growth and development

Organ or target	Effect	Animal class	Reference
Body	↑ Weight	Birds, mammals	(122, 123)
Dorsal fin	↑ Postnatal body growth	Mammals	(124)
	Resorption	Fish	(125)
<i>Larvae</i>			
Gill	↑ Growth	Amphibians	(126)
Tail fin	↑ Growth-via ↑ collagen synthesis		(127, 128)
	↑ Length		(126, 129)
<i>Premetamorphosis</i>			
Hind legs	↓ Growth		(130, 131)
	↓ T ₃ receptor autoinduction		(132)
Tail fin	Prevents loss of tail		(133–134)
	↓ T ₃ receptor autoinduction		(134)
Tail	Regeneration	Reptiles	(93)
Epidermis	Molting	Reptiles, amphibians	(135)
Feather	Growth	Birds	(93)
	Molting		(136–138)
Skin	Metamorphosis-Na channels	Amphibians	(139, 140)
	Incubation patch-defeathering	Birds	(136)
	Incubation patch-epidermal growth	Birds	(141)
	Proliferation of melanocytes	Fish, mammals	(142, 143)
	Proliferation of keratinocytes	Mammals	(144)
	Hair loss for nest building	Mammals (rabbit)	(93)
Hair follicle	Hair growth	Mammals (deer, goat)	(145, 146)
Crop sac	Proliferation of epithelium	Birds	(147, 148)
Fetal lung	Maturation, surfactant production	Mammals	(149–152)
Heart	↑ Ornithine decarboxylase (ODC) activity	Mammals	(153)
Liver	Hepatocyte proliferation	Mammals	(154)
	Activation via		
	Protein kinase C		(155, 156)
	Diacylglycerol		(156)
	MAP kinase		(157)
	Phosphoinositide turnover		(158)
	Induction of growth-related genes		
	IGF-1		(159)
	ODC	Reptiles, mammals	(153, 156, 160, 161)
	β-actin	Mammals	(156)
	c-myc		(156)
	c-fos		(162)
	c-jun		(162)
	c-src		(162)
	↓ Cytokine gene expression in Kupffer cells		(163)
	Induction of growth factors		
	Synlactin		(164–166)
	Liver lactogenic factor		(167)
Liver, kidney	DNA hypomethylation	Mammals	(168)
Kidney	Growth of tubular epithelium	Fish	(169, 170)
	↑ ODC activity	Mammals	(153)
Intestine	↑ Intestinal mucosa	Birds, mammals	(171–173)
	Growth and changes in metabolism		(174, 175)
Muscle	Proliferation of vascular smooth	Mammals	(176, 177)
	Induction of growth-related genes		
	c-myc		(178)
	ODC		(178)
Adipocytes	Preadipocyte differentiation	Mammals	(179)
Pancreas	Proliferation of β-cells	Mammals	(180–183)
	Increased β-cell-to-cell communication		(184–186)
Adrenal	↑ ODC activity	Mammals	(153)
Pituitary	Proliferation of GH3 cells (somatolactotrophs)	Mammals	(187)
Germ cells	↑ Maturation	Mammals	(188, 189)
Gonads	↓ Weight	Birds	(190)
	↑ Weight	Mammals	(191)
	↑ ODC activity		(192)

TABLE 3. Continued

Organ or target	Effect	Animal class	Reference
Prostate	Proliferation of human BPH epithelial cells	Mammals	(193)
	↑ Growth		(194)
	↑ ODC activity		(195)
Seminal vesicle	↑ Growth	Mammals	(196)
Amnion	↑ DNA synthesis and creatine kinase activity	Mammals	(197)
Brain	Astrocyte proliferation	Mammals	(198)
	Tuberoinfundibular hypothalamic development		(199)
Retina	Metamorphosis of visual pigments	Amphibians	(200)
Immune system	↑ Thymus and spleen weights	Mammals	(201)
	Proliferation of lymphocytes		(202, 203)
Tumor growth			
Intestine	Colo-rectal tumor aggressivity	Mammals	(204, 205)
Mammary	Proliferation of	Mammals	
	MXT cells		(206)
	MCF-7 cells		(207)
	T-47D cells		(208)
Lymphoma	Proliferation of malignant B lymphocytes (requires Ig-bound PRL)	Mammals	(209)
	Proliferation due to provirus insertion into PRLR promoter		(210)
	Proliferation of pre-T Nb2 cells		(211)
Promyelocytes	Proliferation of HL60 cell line	Mammals	(212)
Leiomyoma	Associated with ↑ PRL	Mammals	(213)

section of this review, there were some aspects that did not fit well with either the discussion on reproduction or immune function. In liver, in addition to the large number of activities that have already been reported, PRL is also able to increase free intracellular Ca^{2+} concentrations (223), $\text{PGF}_{2\alpha}$ and PGE production (232), and IGF-I production, at least under some circumstances (233).

Finally, many of the multiple functions of PRL in the numerous target tissues that have been identified (Table 1) are enhanced by an up-regulation of PRLRs induced by PRL itself. Thus, PRL is able to sensitize the response of certain tissues by increasing the number of specific PRLRs, representing the first step in the signal transduction process of the hormone (35).

D. Brain and behavior

1. Parental behavior. As shown in Table 5, PRL has been suggested to be involved in parental behavior of fishes, birds, and mammals. In fishes, PRL has been implicated in fish fanning, to provide a constant supply of fresh water to the eggs and to stimulate mucus production, which is used to feed the young after they hatch (108, 236). Another form of parental behavior is foam nest building, in which air bubbles are mixed with mucus to form bubbles during egg laying (237). PRL is also involved in migration. This hormone induces some forms of teleosts to migrate from seawater to fresh water (238) and causes certain forms of salamanders to actively seek and migrate to an aquatic habitat (109). PRL predisposes some species of birds to migrate, as is evidenced by increased nocturnal restlessness and food consumption (239). Birds also respond to PRL by an increase in nesting behavior, nest attendance, and incubation behavior (242–244).

Almost all adult female mammals show some form of maternal care toward their young offspring immediately

after parturition, whereas the response of nulliparous females is less intense or completely absent. The endocrine changes that occur during and at the end of pregnancy, including increased circulating concentrations of PLs and PRL are thought to be responsible for the induction of maternal behavior (245–247). PRL is also considered to be responsible for inducing excessive grooming in rats (248–250). Increased food intake (hyperphagia) can be observed in birds and mammals in response to PRL (190, 241, 251, 252), and in birds, regurgitation feeding has also been shown to be PRL-induced (241, 254).

2. Others. Adaptive stress responses in mammals represent another set of behavioral responses that are induced by PRL (250, 255). Interestingly, PRL has been shown to have analgesic effects (256, 257) that can be mimicked by a number of central nervous system neurotransmitters, including opioids, γ -aminobutyric acid, acetylcholine, and Ca^{2+} and K^{+} channels (258–262). In women, elevated levels of PRL are associated with some psychosomatic reactions, including a form of pseudopregnancy (263). In rats, PRL has been shown to induce increased lordosis behavior (270). Interestingly, the effect does not appear to be associated with rapid electrophysiological action but rather with a neuromodulatory or metabolic effect, since the latency is about 40 min. Elevated circulating PRL is also thought to be responsible for decreased libido (263, 264), increased rapid eye movement sleep (265–267), and an alteration of the sleep-wake cycle (268, 269).

In the hypothalamus, PRL has been suggested to be important for maturation of the neonatal neuroendocrine system (199, 275). In species such as rodents in which the young are born immature, PRL may be provided by maternal milk, which can be absorbed by the intestinal epithelium of neonatal animals for a few days. As much as 20% of ingested PRL is thought to reach the fetal circulation (283). PRL is also

TABLE 4. Endocrinology and metabolism

Organ	Effect	Animal class	Reference
<i>Energy Metabolism</i>			
Brain	Tissue-specific modulation of ATPases	Mammals	(216)
<i>Lipid Metabolism</i>			
Fetal lung	↑ Phospholipid synthesis	Mammals	(149)
Liver	↑ Lipoprotein lipase activity	Mammals	(217)
	↑ Bile secretion		(218)
	↑ Na ⁺ /taurocholate co-transport		(219, 220)
Adipocyte	↑ Lipoprotein lipase activity	Birds	(221)
<i>Carbohydrate Metabolism</i>			
Brain	Modification of enzyme activities	Mammals	(222)
Liver	↑ Glycogen phosphorylase a activity	Mammals	(223)
Pancreas	↑ Insulin secretion	Mammals	(185, 224, 225)
	↓ Glucose threshold for insulin secretion		(226)
	↑ Glucokinase and glucose transporter 2		(227)
<i>Steroid Metabolism</i>			
Adrenal	↑ Steroidogenesis	Mammals	(228, 229)
	↑ Adrenal androgens (DHEA, DHEA-S)		(229)
	↑ Cortisol, aldosterone		(228)
	↑ 21-Hydroxylase activity		(230)
Skin	↑ Steroidogenic enzyme (3βHSD)	Mammals	(231)
<i>Signal Transduction</i>			
Liver	↑ Ca ²⁺ concentration	Mammals	(223)
	↑ PGF _{2α} and PGE		(232)
	↑ IGF-I production		(233)
Many tissues	↑ PRLR number		(35)

DHEA, Dihydroepiandrosterone; DHEA-S, dihydroepiandrosterone sulfate.

known to increase the turnover of dopamine (271–274), to increase the electrical activity of ventral hypothalamic neurons (279), and to augment PKC activity (280). Finally, two effects of PRL have been observed in the retina, the first a decline in the level of TRH receptors (281), and the second, an increase in photoreceptor destruction (282).

E. Reproduction

Actions related to the processes of reproduction represent the largest group of different functions that have been identified for PRL. These are listed in Table 6. The different actions are quite diversified, but certain subcategories can be identified.

1. Nurturing of young. PRL is best known for its actions on the mammary gland. In this complex organ, the growth of the gland that occurs during pregnancy is under the control of a number of trophic factors including estrogen, progesterone, insulin, glucocorticoid, GH, and PRL or PL. The terminal stage of mammary gland development, lobuloalveolar growth, is directly regulated by PRL (93). Although the hormonal requirements for the induction and maintenance of milk production vary in different species, the common factor is that PRL is the hormone primarily responsible for the synthesis of milk proteins (284), lactose (290), and lipids (292), all major components of milk. In addition, PRL has been shown to directly stimulate IGF-I binding protein (296), epidermal growth factor (EGF) (297), a glycosylated mucin (299), parathyroid-like peptide (301), and PRL-inducible proteins (300) in normal and neoplastic mammary tissue.

Although birds do not have a mammary gland, the esoph-

agus of some birds leads to a large chamber known as the crop sac, which connects to the avian stomach. The crop sac serves as an organ for food storage. In pigeons and doves, the crop sac has developed a high degree of sensitivity to PRL, as this hormone is able to stimulate growth of the crop sac, including proliferation and thickening of the epithelium. As the cells move away from the basal layer, they hypertrophy and accumulate fat globules. These cells are finally sloughed off into the crop lumen forming a substance known as crop milk, which is used to feed the young (241). Recently, a number of genes have been shown to be induced in the pigeon crop sac in response to PRL, including annexin Icp 35 (302, 303), lipoprotein lipase (221), ornithine decarboxylase (305), as well as other proteins of unknown function (304, 306).

As has been mentioned above, in some fish, modified skin glands are able to produce mucus, which serves as nourishment for the young. This mucous secretion in many fish is responsive to PRL (108). Thus, as is seen for birds and fish, the production of a substance for the survival of the young is not a feature restricted to mammals.

2. Ovarian actions. In rodents, luteotropic and luteolytic actions of PRL have been recognized for a number of years (307–309). Since progesterone production by the ovaries is necessary for implantation of the fertilized ovum, for maintenance of pregnancy, and for inhibition of ovulation, regulation of progesterone production by the corpus luteum is an important feature in reproduction. In general, the luteotropic action of PRL (338) involves stimulation of progesterone production by luteal cells (341). Progesterone synthesis

TABLE 5. Brain and behavior

Organ or target	Effect	Animal class	Reference
Brain, CNS	Fin fanning; provides fresh water to eggs	Fish	(236)
	Foam nest building		(237)
	Migration	Fish, birds	(238, 239)
	Migration-water drive	Amphibians	(109)
	Nesting behavior	Birds	(136, 240)
	Nest attendance		(136, 241)
	Incubation behavior		(242–244)
	Maternal behavior	Mammals	(245–247)
	Grooming behavior		(248–250)
	Feeding behavior	Birds	(251)
	Hyperphagia	Birds, mammals	(190, 241, 252, 253)
	Regurgitation feeding	Birds	(241, 254)
	Adaptive stress responses	Mammals	(250, 255)
	Induced analgesia mimicking		(256, 257)
	Opioids		(258)
	GABA		(259)
	Acetylcholine		(260)
	Ca ²⁺ channels		(261)
	K ⁺ channels		(262)
	Psychosomatic reaction (pseudopregnancy)		(263)
	↓ Libido		(263, 264)
	↑ REM sleep		(265–267)
	Sleep-wake cycle		(268, 269)
	↑ Lordosis behavior		(270)
Hypothalamus, striatum	↑ Dopamine turnover	Mammals	(271–274)
Hypothalamus	Maturation of neonatal neuroendocrine system	Mammals	(199, 275)
	↓ GnRH secretion		(276, 277)
	↓ Frequency and pulsatility of LH		(278)
	↑ Electrical activity of the VMH neurons		(279)
	↑ PKC activity		(280)
Retina	↓ TRH receptors	Mammals	(281)
	↑ Photoreceptor destruction		(282)

VMH, Ventromedial hypothalamus; REM, rapid eye movement.

is, in turn, affected by activation and inhibition of ovarian steroidogenic enzymes (312–314, 322, 327, 341). Since the maintenance of ovarian progesterone production involves a luteotropic complex rather than a single gonadotropin, PRL certainly acts in concert with LH/hCG and perhaps other factors.

An antagonistic effect of PRL, involving a marked reduction in the weight of the gonads, has been observed in birds (190). In mammals, depending on the stage of the cycle, luteolytic effects of PRL have also been reported (339, 340). Several factors, including PRL, seem to be involved in the destruction of the corpus luteum. In granulosa cells, PRL inhibits estrogen synthesis (319–321) and P450 aromatase (322) and induces α_2 -macroglobulin via activation of Stat5 (336). As a factor regulating the formation and destruction of the corpus luteum, PRL seems to play a major role in modulating the physiological states of estrus, pregnancy, and lactation.

A direct effect of PRL on developmental competence and maturation of oocytes has been reported in rabbits (188, 337). The addition of PRL to the oocyte maturation medium increased the development of organized embryos. In addition, PRL was able to directly inhibit degeneration and decomposition of surface epithelial cells and the disruption of connective tissue at the apex of the follicle wall. PRL inhibited the activity of hCG-stimulated plasminogen activator in ma-

ture follicles. These latter studies suggest that the preovulatory PRL environment can influence oocyte maturation.

3. Uterine actions. In the uterus, PRL is able to increase the level of progesterone receptors, and thus all actions associated with this steroid hormone are enhanced (344, 345). PRL has been reported to induce uterine fluid loss (348) and decrease progesterone metabolism (347). A stimulatory effect is also observed on estrogen receptor levels (349, 350), as well as in conjunction with estrogen, on the general secretory activity of the endometrium (344, 352). PRL promotes blastocyst implantation (355) and increases uteroglobin production (354), leucine aminopeptidase activity (353), and glucose amine synthetase activity (355). Finally, stimulation of prostaglandins and phospholipase A₂ is observed (346).

4. Testicular actions. The physiological role of PRL in males has puzzled investigators ever since the hormone responsible for mammary gland development and lactation in females was shown to be present in the anterior pituitary of males. Initially, no clear function could be ascribed to PRL in male animals or humans (93). More recent data have shown, however, that in general, PRL stimulates testicular functions in most mammals, although as has been reported previously, in birds it causes a decrease in the weight of the gonads (190). In Leydig cells, PRL is involved in the maintenance of cellular morphology (189), increases LH receptor number (191, 356),

TABLE 6. Reproduction

Organ or target	Effect	Animal class	Reference
Mammary gland	Lobuloalveolar growth	Mammals	(93)
	↑ Milk protein synthesis		(284)
	α-Casein		(285)
	β-Casein		(286)
	Whey acidic protein		(287)
	β-Lactoglobulin		(288)
	Late lactation protein of marsupials		(289)
	↑ Lactose synthesis		(290)
	Lactose synthetase		(291)
	Galactosyl transferase		(291)
	α-Lactalbumin		(291)
	↑ Lipid metabolism		(292)
	Acetyl-CoA carboxylase		(293)
	Fatty acid synthase		(294, 295)
	Malic enzyme		(294, 295)
	Lipoprotein lipase		(294, 295)
	↑ IGF-I binding protein		(296)
	↑ EGF		(297)
	↑ 120-kDa protein		(298)
	↑ Muc 1 (glycosylated mucin)		(299)
Crop sac	↑ PRL-inducible protein	Birds	(300)
	↑ Parathyroid-like peptide		(301)
	Growth		(241)
	↑ Mitosis of germinal layer		(147)
	Thickening of epithelium		(147)
	↑ Annexin Icp 35		(302, 303)
	↑ Crop milk polypeptide 58 and 50.5		(304)
	↑ Lipoprotein lipase		(221)
Ovary	↑ Ornithine decarboxylase	Mammals	(305)
	↑ 25-kDa protein		(306)
	Luteotropic and luteolytic actions (dependent on stage of estrous cycle)		(307–309)
	Ovum maturation		(310)
	↓ Folliculogenesis		(311)
	↓ 3β-HSD		(312)
	↓ Aromatase		(313)
	Potentiate effects of LH on 3β-HSD		(314)
	↓ Ovulation		(315)
	↓ Estradiol and induce ovarian regression	Birds	(316)
Granulosa cells	↓ Plasmin generation in preovulatory follicles	Mammals	(317, 318)
	↓ Estrogen production	Mammals	(319–321)
	↓ P450 aromatase		(322)
	↑ Use of extracellular lipoproteins		(323, 324)
	↑ Progesterone production		(320, 324, 325)
	↑ LH receptors		(326)
	↑ 20α-OH progesterone (humans)		(327)
	↓ Progestins (humans)		(328–330)
	↓ Luteinization		(331)
	↓ cAMP and steroidogenesis (rats)		(332)
	Counteract morphological effects of LH		(333)
	↑ Progesterone in cocultures of splenic macrophages removed at proestrus		(334)
	↑ DAG		(335)
	↑ 101 ₂ -Macroglobulin		(336)
Oocytes	↑ Developmental competence and maturation (rabbits)	Mammals	(188, 337)
Luteal cells	Luteotropic action	Mammals	(338)
	Luteolysis in pregnancy (marsupials)		(339)
	Luteolysis during estrous cycle		(340)
	↓ 20α-HSD		(341)
	↑ Progesterone		(307)
	Control of delayed implantation and steroidogenesis (bats)		(342)
	↓ 37-kDa protein		(343)

TABLE 6. Continued

Organ or target	Effect	Animal class	Reference
Uterus	<ul style="list-style-type: none"> ↑ Progesterone receptors and progesterone effects ↑ PGE₂, phospholipase A₂, prostaglandin G/H synthase ↑ Fluid loss ↓ Progesterone metabolism ↑ Estrogen receptors ↓ Myometrial contraction ↑ General secretory activity of endometrium (with estrogen) ↑ Leucine aminopeptidase activity (with estrogen) ↑ Uteroglobin production Promote blastocyst implantation ↑ Glucose amine synthetase activity 	Mammals	(344, 345) (346) (347) (348) (349, 350) (351) (344, 352) (353) (354) (355) (355)
Testis	In general, ↑ activities	Mammals	
Leydig cells	<ul style="list-style-type: none"> Involved in maintenance of cell morphology ↑ LH receptors ↓ Aromatase activity (with LH) ↑ Steroidogenesis and androgen production (cooperation with LH) 	Mammals	(189) (191, 356) (357) (356, 358–360) (191)
Sertoli cells	↑ FSH receptors	Mammals	(361)
Germ cells	↑ Total lipids	Mammals	(362)
	↑ Spermatocyte-spermatid conversion		(189)
Spermatozoa	<ul style="list-style-type: none"> ↑ Ca²⁺ binding and/or transport of ejaculated and epididymal spermatozoa Energy metabolism: ↑ ATPase activity ↑ Fructose rate ↑ Glucose oxidation ↓ Zn²⁺ content Maintenance of mobility and attachment to oocyte Shortening optimal preincubation period to acquire capacitation 	Mammals	(363) (364) (365) (366) (364, 367) (368) (369) (369)
Epididymis	<ul style="list-style-type: none"> Energy metabolism: ↑ Glycogenolysis and hexophosphate enzymes ↑ Sialic acid ↑ β-galactosidase and α-mannosidase activities ↑ Lipids ↓ Glycoprotein metabolism 	Mammals	(370) (371) (372) (373) (374)
Seminal vesicle	<ul style="list-style-type: none"> Hyperprolactinemia: ↓ sperm fertilizing and mobility capacity Hypoprolactinemia: azoospermia ↑ Fluid lipids ↑ Lipogenesis ↑ Phosphomonoesterase and acid phosphatase ↓ Glycosylation 	Mammals	(375, 376) (377) (378) (379) (380) (381)
Prostate	<ul style="list-style-type: none"> ↑ Weight ↑ Nuclear uptake of DHT ↑ Androgen receptor Involvement in estrogen-induced inflammation ↑ Epithelial secretory function ↑ Energy metabolism: Monosaccharide formation Amino acid oxidation and transamination ↑ Ornithine decarboxylase ↑ Citric acid secretion: ↑ Mitochondrial aspartate Aminotransferase (via PKC) Citrate oxidation and m-asconitase: Ventral prostate: ↑ Dorsal prostate: ↓ ↑ Pyruvate dehydrogenase E1α ↑ Aspartate transporter ↑ C3 subunit of prostatein, probasin, RWB gene ↑ IGF-1 and IGF-1 receptor 	Mammals	(123) (382–384) (385, 386) (387) (381, 388) (389) (390) (195) (391) (392) (393) (394) (391) (395) (386)

HSD, Hydroxysteroid dehydrogenase; DAG, diacylglycerol; DHT, dihydrotestosterone.

TABLE 7. Immunoregulation and protection

Organ or target	Effect	Animal class	Reference
Spleen	↑ Weight	Mammals	(201)
Thymus	↑ Weight	Mammals	(201)
	↑ Thymulin production		(397)
Submandibular gland	↑ Immunostimulatory activity	Mammals	(398)
Lymphocytes	↑ Hormonal immunity	Mammals	(399)
	↑ Adjuvant arthritis response		(399-401)
	↑ Cellular immunity		(402)
	↑ Antibody formation to sheep RBC		(403)
	↑ IgG and IgM antibodies		(404-406)
	Reverse hypophysectomy-induced anemia, leukopenia, and thrombocytopenia		(407)
	↑ Proliferation		(203, 408-410)
	↑ IL-2 receptors		(203, 411)
	↑ EPO receptors		(412)
	↑ PRL receptors		(413, 414)
	↓ Apoptosis		(415, 416)
	↑ IFN- γ		(417)
	↑ Graft rejection		(418)
	↑ <i>c-myc</i>		(201)
	↑ DNA synthesis		(201)
	↑ T cell engraftment		(419)
	↑ DNA synthesis	Birds	(420)
Nb2 cells	↑ Proliferation	Mammals	(211)
	↓ Apoptosis		(416, 421)
	Genes induced:		
	IRF-1		(422, 423)
	<i>c-myc</i>		(424)
	<i>c-fos</i>		(424)
	Ornithine decarboxylase (ODC)		(425)
	hsp 70		(426)
	β -Actin		(427)
	<i>pim-1</i>		(428)
	<i>gfi-1</i>		(429)
	<i>bcl-2</i>		(430)
	<i>bax</i>		(430)
	T-cell receptor γ -chain		(431)
	Cyclins D2 and D3		(432)
	Cyclin E, <i>cdk2</i> , <i>cdk5</i> , E2F-1		(433)
	Clone 15, nuclear movement protein		(434)
	GnRH		(435)
	GnRH receptor		(435)
	Proteins activated:		
	Jak2		(436-439)
	Fyn		(440)
	Stat proteins		(436, 441, 442)
	Ras		(443, 444)
	Raf		(445)
	Vav		(446)
	Grb2		(444)
	Sos		(444)
	Shc		(444)
	MAP kinase		(443, 447)
	Stathmin		(448, 449)
	IRS-1		(450)
	PTP-1D (SHP-2)		(451)
	PKC		(452, 453)
	Casein kinase II		(454)
	PTK		(452)
	PI3 kinase		(450, 455)
	<i>cbl</i>		(456)
	S6 kinase		(457)
	G proteins		(458, 459)
	PLC		(460)
	Amiloride-sensitive Na ⁺ /H ⁺ exchange system		(461)
NK cells	↑ Susceptibility of primary leukemia cells	Mammals	(462)
	↑ Cytotoxic effects		(463)
	↑ DNA synthesis		(464)

TABLE 7. Continued

Organ or target	Effect	Animal class	Reference
Macrophages	↑ Activation	Mammals	(408)
	↑ Cytokine gene expression in Kupffer cells following hemorrhage		(163)
	↓ Monoblastic growth-synergy with IFN- γ		(465)
	↑ Superoxide anion responsible for killing pathogenic organisms		(466, 467)
	↑ Nitric oxide and protection against bacterial infection		(468)
Polymorphonuclear cells	↓ Direct and spontaneous migration	Mammals	(469)
Thymic nurse cell complexes	Regulation of lymphocyte/epithelial cell adhesive interactions	Mammals	(470)
Mammary gland	↑ IgA-secreting plasma cells	Mammals	(471)
Liver	Induction of coagulation factor XII	Mammals	(472)
Skin	Melanogenesis	Fish	(142)
Pathology			
Lymphocytes	Treatment for immunodepression following severe hemorrhage	Mammals	(473)
	↑ Autoimmune responses		
	Systemic lupus erythematosus		(405, 474, 475)
	Acute experimental allergic encephalomyelitis		(476)
	Rheumatoid arthritis		(477)
	Adjuvant arthritis		(399, 400, 478)
	Graft <i>vs.</i> host disease		(479)

and along with LH, decreases aromatase activity (357) and increases steroidogenesis and androgen production (356, 359, 360). In Sertoli cells, PRL has been shown to increase FSH receptor numbers (361). In germ cells, PRL increases total lipids (362) and increases the spermatocyte-spermid conversion (189). Several effects on spermatozoa have been reported, including an increase of calcium binding and/or transport of ejaculated and epidymal spermatozoa (363) as well as an increase in energy metabolism (364), a maintenance of mobility and attachment to the oocyte (369), and a reduction in the time required to achieve capacitation (369).

5. Male sex accessory actions. In rats, effects on accessory sex organs are well established. In addition to increasing the weight of the prostate and seminal vesicle, in conjunction with androgens (194, 196), PRL also has metabolic effects on sex accessory organs. In the epididymis, energy metabolism is increased (370) and, in addition, sialic acid and lipid levels are augmented (371, 373). In the seminal vesicle, PRL increases lipid concentrations in the fluid (378), lipogenesis (379), and phosphomonoesterase and acid phosphatase activities (381). The effects of PRL on prostate include increased levels of androgen receptor (385, 386), involvement in estrogen-induced inflammation (387), increased epithelial secretory function (381, 388), augmented energy metabolism, and an enhancement of citric acid production (391). PRL has also been reported to stimulate the level of IGF-I and IGF-I receptor in the prostate (386) and to increase the production of other prostate-specific proteins (395).

F. Immunoregulation and protection

Although actions of PRL on the hematopoietic system were suggested in early studies involving treatment of hypophysectomized rats with PRL (396), it was not until much later that a clear role of PRL in the immune system was established. Table 7 summarizes the multiple effects of PRL in the regulation of immune function. Injection of PRL into hypophysectomized rats causes an increase in the weight of

the spleen and thymus (201). In addition, PRL activates an immunostimulatory action of the submandibular gland (398) and augments the production of a thymic hormone, thymulin (397).

In lymphocytes, PRL is known to increase hormonal and cellular immunity (399, 402), to reverse anemia, leukopenia, and thrombocytopenia induced by hypophysectomy (407), to increase antibody formation, including IgG and IgM antibodies (404–406), and to induce cellular proliferation (203, 408–410), although this last function is somewhat controversial, as not all investigators are able to reproduce PRL-induced proliferation of lymphocytes. Recently, in fact, it has been suggested that the proliferative action of PRL of recombinant origin and even natural preparations may be due to contamination with endotoxins, which are responsible for the immunostimulatory actions that have been observed (480). This interesting observation awaits confirmation. Along with effects on proliferation, PRL has also been shown to increase DNA synthesis, and *c-myc* expression (201).

PRL has also been reported to increase receptor levels for interleukin (IL)-2 (203, 411), EPO (412), and PRL (413, 414). Recently, in addition to stimulating proliferation, PRL has been shown to inhibit apoptosis of lymphocytes (415, 416)(see below). Administration of PRL is also associated with increased graft rejection (418, 481) and an increase in T cell engraftment (419). In natural killer cells, PRL has been reported to increase DNA synthesis (464) and augment cytotoxic effects (463), as well as increasing susceptibility of primary leukemic cells (462).

Activation of macrophages was originally thought to be an action of GH. More recently, however, it has been shown that macrophage activation (408) and superoxide anion production responsible for killing pathogenic organisms (466, 467) are effects mediated by the PRLR. The protective effect against *Salmonella* infection appears to be mediated by nitric oxide production (468). In synergy with IFN- γ , PRL decreases monoblastic growth (163) and increases cytokine gene expression in Kupffer cells after hemorrhage (163).

PRL reduces direct and spontaneous migration (469) of polymorphonuclear cells (469), regulates lymphocyte-epithelial cell adhesive interactions in the thymic nurse cell complex (470), and augments IgA-secreting plasma cells in mammary gland (471).

In terms of more specific protective effects, PRL induces the production of coagulation factor XII by the liver (472) and, in fish, regulates melanogenesis (144).

Finally, as listed in Table 7, multiple functions of PRL are observed in the pre-T lymphoma Nb2 (for review, see Ref. 482). These cells respond to very low concentrations of PRL by a marked proliferation (211) and, in fact, are used routinely as a bioassay of PRL (483). Within the past few years, studies on Nb2 cells have also highlighted the antiapoptotic properties of PRL on lymphoid cells. This effect appears mediated by gene products known to be involved in apoptosis, such as Bcl-2 and Bax (416, 430), although involvement of newly identified proteins such as *pim-1* (428) or *Bag-1* (484) has also been proposed. Almost all the activities reported for Nb2 cells in Table 7 are associated with the mechanism of action of PRL: these include the induction of many growth-related genes, and the activation of a number of proteins implicated in the signal transduction process. These effects are discussed in detail in Section VI.

G. Actions associated with pathological disease states

In humans, hyperprolactinemia has been shown to be associated with amenorrhea, galactorrhea, and impotence (485). The inhibitory effects on the reproductive processes may be due to both central and peripheral actions of PRL. In some women, elevated PRL is associated with a psychosomatic state of pseudopregnancy (263, 264).

Interestingly, however, no genetic diseases associated with a mutation of the gene encoding PRL or the PRLR have been identified in humans or animals. Either these genes are not important or they are essential to the proper survival of the species. As described below, knockout of the PRLR gene in mice is not lethal but does produce major reproductive defects in females, which would, of course, affect reproductive function and survival.

An excessive volume of amniotic fluid, known as polyhydramnios, is associated with decreased levels of amniotic fluid PRL (120) or PRLR levels in the chorion laeve (486). This effect may be related to the osmoregulatory role of PRL during fetal life and to the inhibitory effect on amniotic fluid volume observed in monkeys (119).

PRL has been associated with a number of different forms of cancer. For example, PRL is thought to increase colorectal tumor aggressivity (204, 205), induce the proliferation of several lines of human breast cancer (206–208), activate malignant B lymphocytes (209) and lymphoma cells (211), and induce the proliferation of promyelocytes (212). Benign fibromuscular myometrial tumors (leiomyomas) have been shown to produce more PRL than control myometrium (213); thus, locally produced PRL may exert a mitogenic action on the growth of these tumors.

PRL has been shown to be increased and to effect a number of autoimmune states, such as systemic lupus erythematosus (405, 474, 475), acute experimental allergic encephalomyelitis

(476), rheumatoid arthritis (477), adjuvant arthritis (478), and graft vs. host disease (479), e.g., as a marker of rejection in heart transplantation (481). PRL has also been suggested to be involved in the etiology of cystic fibrosis (121), although the precise mechanism remains unclear.

VI. Signal Transduction by the PRLR: Structure-Function Relationships

All of the actions described above result from the interaction of PRL with its receptor in various target cells, which leads to the activation of a cascade of intracellular events. In this section, we summarize the current state of knowledge concerning signal transduction of the PRLR. As indicated previously, Nb2 cells are one of the preferred models with which to study PRL actions, and a list of genes induced and proteins activated by lactogenic stimulation of these cells can be found in Table 7.

A. The JAK-Stat pathway

1. Janus kinases.

JAK family. The cytoplasmic tail of the PRLR, whatever the isoform, is devoid of any consensus sequence for enzymatic activity, including kinase activity (35, 487), as are all cytokine receptors identified thus far. However, hormonal stimulation of the PRLR leads to tyrosine phosphorylation of several cellular proteins, including the receptor itself (488), and tyrosine kinase inhibitors were shown to inhibit the mitogenic and anabolic effects of lactogenic hormones in lymphoid Nb2 cells (489). Until 4 yr ago, this observation was unsolved. In the early 1990's, Wilks and colleagues (490, 491) identified, by low-stringency hybridization and PCR approaches, a new family of protein tyrosine kinases to which they gave the acronym JAK (for just another kinase). Regarding the presence of two kinase-like domains in these kinases, the name JAK was also proposed as an acronym for Janus kinase, in reference to the dual-faced Roman god Janus (492). To date, the JAK family includes four members, termed JAK1, JAK2, JAK3, and Tyk2 (reviewed in Ref. 493). All cytokine receptors work in combination with one or several JAKs to transmit the hormonal signal within the cell (41, 44, 45, 493–495). In 1994, as demonstrated 1 yr before for the GHR (496), JAK2 was identified by three laboratories as the JAK kinase associated with the PRLR (438, 439, 497). Since this major discovery, important steps in understanding the complex field of signal transduction by the PRLR have been made and are summarized in the following sections. A schematic representation of the current knowledge of the main PRLR-activated signaling pathways is proposed in Fig. 3.

Association of JAK2 with the PRLR. Although involvement of JAK1 has also been proposed in the particular context of mouse lymphoid BAF/3 cells transfected with the PRLR (498), JAK2 is unambiguously the major PRLR-associated Janus kinase. Using mutant cell lines defective in JAK1 or JAK2, Stark and colleagues (499) have very recently confirmed this assumption and showed that although milk protein gene induction required the latter, the former was dispensable. Possible interactions between JAK2 and JAK3 in avian PRLR signaling have been suggested, but await further




FIG. 3. Schematic representation of the PRLR signaling pathways. Long and short isoforms of rat PRLR are represented. PRLR activates Stat1, Stat3, and, mainly, Stat5. Interaction of Stat5 with the glucocorticoid receptor (GR) has been reported. Whether the short PRLR isoform activates the Stat pathway is currently unknown. PRAP (PRLR-associated protein) seems to interact preferentially with the short PRLR. The MAP kinase pathway involves the Shc, Grb2, Sos, Ras, Raf cascade and is presumably activated by both PRLR isoforms. Connections between the JAK-Stat and MAPK pathways have been suggested. Interactions between receptors and Src kinases (e.g., Fyn), SHP2, IRS-1, PI-3 kinase, and other transducing molecules remain unclear.

confirmation (59). JAK2 is constitutively associated with the PRLR (439, 500), i.e., it is not induced by ligand binding, contrary to what has been observed for the GHR (496). The PRLR-JAK2 interaction involves the membrane-proximal region of the PRLR cytoplasmic domain, in agreement with the ability of the short PRLR isoform to associate with the kinase (501). This region includes the consensus box 1, which is highly enriched in proline residues (I²⁴³-P²⁴⁴-P²⁴⁵-V²⁴⁶-P²⁴⁸-G²⁴⁹-P²⁵⁰ in rat PRLR) and is believed to adopt the typical folding of SH3-binding domains (502). Alanine substitutions of individual residues within box 1 of the rat PRLR have shown that the most C-terminal proline (P²⁵⁰) is critical for association with and subsequent activation of JAK2 (503). However, sequence analysis of Janus kinases failed to identify any consensus SH3 motif. Current knowledge thus suggests either that an adapter links JAK2 and the PRLR, by interacting with the PRLR box 1 through an SH3 domain and with JAK2 through another mechanism, or that the receptor-kinase interaction involves a mechanism different from the known SH3-SH3 binding domain. In agreement with the latter hypothesis, O'Neal and colleagues (504) recently proposed that peptides including the amino acid sequence of PRLR box 1 adopt a folding different from that of classic SH3-binding domains and, therefore, are not able to interact with protein SH3 domains. These authors also suggested that

isomerization of the second proline of the consensus P-x-P motif might regulate the activation of the receptor in a "on/off" switch manner (504). Although this proposal relies on observations made in the particular context of spectrometric analysis (circular dichroism, nuclear magnetic resonance) of peptides, it is in good agreement with previously reported mutational studies emphasizing the functional importance of the second proline residue of the P-x-P motif in both the PRLR and the GHR (503, 505-507). Finally, although these data clearly demonstrate that the membrane-proximal box 1 is an absolute requirement for associating with the kinase JAK2, involvement of additional residues toward the C terminus cannot be ruled out. In this respect, that the wild-type rat short PRLR (291 aa) associates with JAK2 in fibroblasts transfected with a plasmid encoding this receptor (501) whereas a mutant form of the long PRLR truncated 11 residues upstream (analog named T280, containing the 280 N-terminal aa) does not (437, 508), might reflect some subtle involvement of the C-terminal tail of the short PRLR in JAK2 binding (see below).

To the best of our knowledge, no mutational study aimed at mapping the region of JAK2 interacting with the PRLR has yet been reported. However, it was recently shown that the N-terminal fifth of JAK2 is involved in GHR binding (509); whether this region binds directly to the GHR is unknown.

Activation of JAK2 by the PRLR. It is usually assumed that activation of Janus kinases occurs by transphosphorylation of tyrosines upon ligand-induced oligomerization of cytokine receptors, which brings two JAK molecules close to each other (44, 495). Based on JAK kinase activation by chimeric receptors in which various ECDs of cytokine or tyrosine kinase receptors were fused to the IL-2 receptor β -chain, we have recently suggested that the stoichiometry of oligomerized cytokine receptors, namely homodimerization, might be involved in the specific recruitment of JAK2 by the PRLR (509a). Identification of tyrosine residues of JAK2 that are phosphorylated after PRLR activation remains to be performed. As expected, PRLR mutants unable to associate with JAK2, such as a box 1-deleted PRLR (501) or the C-terminal-truncated form T280 (437), are also unable to induce tyrosine phosphorylation of the kinase. An elegant experiment using chimeras of GM-CSF/PRL receptors (extracellular of GM-CSFR α and β , intracellular of PRLR) recently showed that heterodimerization of the short and the intermediate PRLR cytoplasmic tails achieves inactive complexes unable to stimulate JAK2 autophosphorylation (510), whereas both have been reported to associate with and activate the kinase in the context of their respective wild-type receptor (501). These data, suggesting that two copies of box 1 alone are insufficient for achieving JAK2 activation (510), are in agreement with results obtained with the T280 truncated long PRLR (437) but contradict the ability of short PRLR to activate JAK2 (501). These discrepancies may result from structural disturbance in the particular context of modified receptors (deleted or chimeric), but could also emphasize the functional significance of the 30 C-terminal residues of the short rat PRLR (aa 262–291; Fig. 2A), which differ from the equivalent region in the intermediate and long PRLR isoforms (35) and, thereby, might confer distinct properties to the receptor whether homodimerization or heterodimerization occurs. Based on further analysis of GM-CSFR/PRLR chimeras, Clevenger and colleagues (510a) recently showed that heterodimerization of the wild-type cytoplasmic domain of the intermediate PRLR with any truncated form of this PRLR, even including box 1, was detrimental to JAK2 autophosphorylation. They also observed that mutation of both tyrosines 309 and 382 within a single PRLR chain prevents JAK2 activation, which strengthens the belief that juxtaposition of strictly identical cytoplasmic domains is required for proper activation of the kinase. In this respect, we have recently shown, using several cell lines [Chinese hamster ovary (CHO) cells, 293 human fibroblasts, bovine mammary gland epithelial (BMGE) cells], that the short PRLR functions as a dominant negative isoform, inhibiting the activation of milk protein gene transcription by the receptor complex through heterodimerization. Indeed, in human embryonic kidney 293 fibroblasts cotransfected with cDNAs encoding the short and long PRLR isoforms, such heterodimers are shown to occur and result in an absence of JAK2 activation (511, 512). All these observations are in good agreement with our proposal that the homodimeric stoichiometry of some cytokine receptors is one of the major features leading to activation of JAK2 (509a).

Activation of JAK2 by the PRLR occurs very rapidly after hormonal stimulation (within 1 min, see Ref. 501). This sug-

gests that the Janus kinase occupies a central and very upstream role in the activation of several signaling pathways of the PRLR. Accordingly, PRLR mutants that fail to associate with and/or to activate JAK2 were all reported inactive in any of the bioassays performed (437, 499, 501, 503, 510). When activated, JAK2 phosphorylates tyrosine residues on different target proteins, the best identified of which are the receptor itself (next paragraph) and a family of transducing proteins termed Stat (signal transducer and activator of transcription) (see Section VI.A3).

2. PRLR phosphorylation. Although the three PRLR isoforms are able to activate JAK2 (see above), the short isoform does not undergo tyrosine phosphorylation (501) in spite of the presence of four tyrosines in its cytoplasmic domain (Fig. 2A). Since phosphotyrosines are potential binding sites for transducer molecules containing SH2 (or any other phosphotyrosine recognizing) domains, particular attention is usually devoted to these amino acids, which play a central role in receptor signaling. In the intermediate PRLR isoform, which contains only three tyrosine residues in its cytoplasmic domain, we have identified the most C-terminal (Tyr 382) as the sole tyrosine undergoing phosphorylation upon receptor/JAK2 activation (513). Substitution of the equivalent tyrosine (Tyr 580) for a phenylalanine in the long isoform does not entirely abolish receptor tyrosine phosphorylation, however, indicative of the presence of at least one other potential phosphorylation site among the eight remaining tyrosines (513). To determine these potential phosphorylation sites, we recently performed a selected mutagenesis of each tyrosine of the rat long PRLR (514). Our results indicate that, in addition to Tyr 580, PRL is able to stimulate phosphorylation of tyrosines 473 and 479 in the basal state, but that in cells overexpressing JAK2, tyrosines 309, 402, and 515 can be phosphorylated in the particular context of PRLR containing a single tyrosine residue (Fig. 2B). These data can be correlated with those obtained from similar studies performed on the GHR, which is homologous to the long PRLR and also contains several tyrosine residues (nine for the rabbit GHR, eight for porcine GHR). Using point-mutated, domain-deleted, or C-terminal-truncated GHR analogs, several tyrosine residues have been mapped as potential phosphorylation sites (515–521). Taken together, these experiments performed on two closely related receptors indicate that tyrosine residues able to undergo phosphorylation are numerous and may be redundant, as illustrated by the intermediate PRLR isoform, which contains a single phosphorylation site and, so far, has been shown functionally equivalent to the long PRLR isoform (513, 522). To the best of our knowledge, no correlation between tyrosines that are preferentially phosphorylated by JAK kinases and their surrounding amino acids has been identified.

Despite the potential phosphorylation of several tyrosine residues within the long PRLR, the most C-terminal (Tyr 580) appears critical for PRLR functional activity since a PRLR mutant containing this sole tyrosine can mimic the wild-type receptor in stimulating reporter genes containing PRL-inducible sequences such as the β -casein promoter or Stat5 DNA-binding sites, while any PRLR analog containing one of the other single tyrosine displays drastically reduced or

even no activity in such bioassay (514). Furthermore, we have shown that the presence of the C-terminal tyrosine on both monomers of the dimerized PRLR complex is required to achieve a fully functional receptor on a PRL-responsive reporter gene (512). In good agreement, Clevenger and colleagues reported recently that mutation of Tyr 382 within the intracellular domain of a single PRLR (intermediate form) involved in a dimerized complex strongly decreases PRL-induced proliferation of BA/F3 cells (510a). In contrast, however, point-mutation of Tyr 580 in the long PRLR isoform has only a partial effect on PRLR function (513), whereas the absence of this residue in bovine and cervine tissues that naturally lack this C-terminal tyrosine compared with other species is not detrimental to receptor activity (523, 524), indicating that alternative intracellular tyrosines can be used. For example, a recent study has suggested an important role of Tyr 309 in the transcriptional activation of the IRF-1 promoter (507), most probably through Stat1 activation (see below). Similarly, dimerization of GM-CSFR/PRLR chimeras in which a single cytoplasmic domain lacks Tyr 309 only achieves partial activation of JAK2 and Fyn kinases and reduced ligand-induced proliferation of BA/F3 cells (510a). Tyrosine 309, which in our bioassays was not critical for β -casein gene activation (513), might thus be required for some biological functions and/or in some cell types.

Why is receptor tyrosine phosphorylation required? Although available data remain limited, it appears that PRLR-mediated cell proliferation does not necessarily require tyrosine phosphorylation of the receptor. For example, a mutant form of the long PRLR truncated after 94 cytoplasmic residues (G328) maintains the ability to activate Stats 1, 3, and 5 and to transduce a significant proliferative signal in murine 32D cells, presumably through signaling pathways involving these Stat molecules (see below) (437). Although the phosphorylation status of this G328 mutant was not established by the authors (437), a mutational study of tyrosines in the intermediate PRLR (513) suggests that this G328 mutant is not phosphorylated, although phosphorylation of tyrosine 309 cannot be totally discarded (514). These results are in agreement with the observation that the nonphosphorylated short PRLR can induce proliferation of NIH-3T3 fibroblasts (525). However, some contradiction exists since the short PRLR was otherwise reported to be unable to stimulate the proliferation of FDC-P1 and Ba/F3 cells (526), as is a homodimerized intermediate PRLR truncated at residue 322 (thus containing a longer tail than the short isoform) expressed in Ba/F3 cells (510a). Such discrepancies might reflect some differences inherent to the cell types (*e.g.*, fibroblastic *vs.* hematopoietic) that were used for these experiments. It thus appears premature to deduce any general rule with respect to PRLR domains required for proliferation. Beyond receptor phosphorylation, it is particularly interesting to note that heterodimerization of wild-type rat PRLR cytoplasmic domains (short/long, short/intermediate, long/intermediate) fails to induce BA/F3 cell proliferation (510a). Since tissues expressing the PRLR presumably express more than a single isoform, this observation might be of considerable functional significance and represents an open area of investigation.

In contrast to cell proliferation, transactivation of reporter

genes (luciferase/chloramphenicol acetyltransferase) controlled by promoters of PRL target genes (*e.g.*, milk proteins such as β -casein) requires additional C-terminal region(s) of the receptors, most probably for the presence of phosphorylated tyrosines within these domains. For example, the rat short PRLR (501), as well as the nonphosphorylated Nb2 mutant lacking the C-terminal tyrosine 382 (513), are unable to stimulate the β -casein promoter and both function as dominant negatives of the long form, strengthening the likelihood that functional complexes require a phosphorylated tyrosine on both receptors forming the homodimer (512). Activation of the IRF-1 promoter, a growth-related PRLR-target gene (441, 527), depends on both tyrosines 309 and 382 of the intermediate PRLR, and it has been suggested that these tyrosines, when phosphorylated, may serve as docking sites for recruiting Stat1 (482, 507). However, we have shown that a mutant form of the long PRLR deleted of all its cytoplasmic tyrosines maintains the ability to transmit a (weak) lactogenic signal in the particular context of JAK2 overexpression, suggesting that the kinase can partially initiate PRLR signaling pathways (514). In agreement, it has been suggested that JAKs by themselves are capable of generating a signal and activating promoters in the absence of a specific ligand-receptor interaction, correlating the ability of the kinase to phosphorylate Stat5 *in vitro* (528).

In conclusion, although current data do not unambiguously establish receptor phosphotyrosines as an absolute requirement for the activation of PRLR target genes, they undoubtedly enhance the signal of PRL on these genes.

3. Stat proteins.

Stat family. Stat is a family of latent cytoplasmic proteins of ~90–100 kDa identified within the past 4 yr on the basis of their involvement in cytokine receptor signaling. The Stat gene family currently contains eight members: Stat1 (α and β), Stat2, Stat3, Stat4, Stat5a, Stat5b, Stat6 (or IL-4 Stat), and dStat, a Stat5 homolog found in *Drosophila*. Stats contain five conserved features: a DNA-binding domain, a SH3-like domain, a SH2 domain, a ubiquitous tyrosine and a C-terminal transactivating domain (from the N- to C-terminus, respectively). A consensus model of Stat activation has been proposed on the basis of data collected from studies of the different cytokine receptors (for additional information, see previously published reviews on Stat proteins, Refs. 44, 495, 529–532). First, the cytokine-bound receptor undergoes tyrosine phosphorylation by the associated Janus kinase. Second, the phosphorylated tyrosine interacts with the SH2 domain of a Stat, making the latter a part of the receptor-JAK complex. Third, the receptor-bound Stat is phosphorylated by the Janus kinases belonging to the complex. Fourth, the phosphorylated Stat dissociates from the receptor, homo- or heterodimerizes through an interaction involving the phosphotyrosine of each monomer and the SH2 domain of the other Stat molecule, and finally the dimer translocates to the nucleus where it activates specific DNA elements found in the promoters of cytokine target genes.

Stats involved in PRLR signaling. The specificity of a cytokine receptor is believed to be driven, at least in part, by the number and type of Stat proteins it can activate. Three mem-

bers of the Stat family have been thus far identified as transducer molecules of the PRLR: Stat1, Stat3 and, mainly, Stat5.

In 1994, the transcription factor named MGF (mammary gland factor) was identified from sheep mammary gland (533) and, in view of the high structural similarity with the other Stat proteins known at that time, it was further renamed Stat5 (528, 530). Within the past 3 yr, Stat5 cloning from mouse (534–536), rat (537), and human (538, 539) revealed the existence of two genes (Stat5a and Stat5b) encoding several isoforms that show 90–95% similarity, the major differences lying within the C-terminal domain. All isoforms possess the functionally essential tyrosine 694, identified by Gouilleux and collaborators (528) as the tyrosine being phosphorylated by JAK2. In agreement, PRLR mutants not able to associate/activate Janus kinases are unable to activate Stat5 (508, 514). Similarly, when the C-terminal domain of Stat5 is truncated (which removes the tyrosine phosphorylation site), such mutants function as a dominant-negative (540–542). Finally, it is noteworthy that in the specific context of cytokine chimeric receptors, JAK1 and possibly JAK3 can also phosphorylate Stat5 on tyrosine (543).

In agreement with the consensus model of Stat recruitment by receptor phosphotyrosines, we have shown that in the long PRLR, tyrosines 580, 479, and 473 (which are all phosphorylated in the context of single-tyrosine mutants), are sufficient to activate Stat5, although Tyr 580 is clearly the most potent in this respect (514). Similarly, mutation of the C-terminal tyrosine 382 in the intermediate PRLR, which abolishes tyrosine phosphorylation of the receptor (see above), prevents PRLR-mediated activation of the β -casein promoter (513). These data strongly suggest that in both long and intermediate PRLR isoforms, this particular tyrosine residue is involved in Stat5 recruitment and/or activation (Fig. 2B). In contrast, others have reported that a C-terminal truncated form of the long PRLR (G328, lacking Tyr 382) is able to stimulate Stat5 tyrosine phosphorylation (508), although the transcriptional activity of this mutant on a PRL-responsive reporter gene was not assessed. Others have shown that the C-terminal tail of the rabbit PRLR is not an absolute requirement, but considerably amplifies the transcriptional activation of PRLR-sensitive genes such as the β -lactoglobulin (500). Although these apparent discrepancies might reflect species specificity (of cellular systems used for bioassays, or of PRLR), they might also indicate that Stat5 tyrosine phosphorylation is required (528), but not necessarily sufficient for activation of transcriptional activity. Accordingly, it has been shown recently that serine/threonine phosphorylation is an absolute requirement for transcriptional activation of Stat5 by the IL-2R (544). Interestingly, Stat1, Stat3, Stat4, and sheep Stat5 contain potential MAP kinase serine phosphorylation sites (P-x-S-P) in their C termini (545). In human, mouse, and rat Stat5a, but not Stat5b, a very similar tetrapeptide is found at the homologous position (R-L-S-P). Although this sequence does not perfectly match the MAP kinase consensus site, it might be a target for another proline-directed kinase (539). Beadling and colleagues (544) have shown that IL-2-induced activation of Stat5 is not mediated by the ERK2/MAP kinase pathway. PKC α and casein kinase II have been proposed as candidates for serine phosphorylation of Stat5 (537, 544). Such a hy-

pothesis may correlate with the previously reported implication of both kinases in the regulation of the β -casein gene by PRL (546, 547). Although functional distinction between Stat5a and Stat5b remains to be investigated in detail, it might be partly correlated with these putative serine/threonine phosphorylation sites. In this respect, heterodimerization of Stat5a and Stat5b has been recently reported by several investigators (548–550). Moreover, Kirken *et al.* (550) have recently shown first that PRL induced phosphorylation of both Stat5a/b on serine, but not on threonine in Nb2 cells, and second that the kinetics of this phosphorylation were markedly different for both Stats (550). Whether this observation reflects any functional difference between these two closely related transcription factors awaits further investigation. Finally, Yu-Lee and colleagues (482, 507, 551) have shown recently that Stat5a and Stat5b exert an inhibitory effect on PRL-inducible IRF-1 promoter activity, and these authors have proposed this inhibition to involve sequestration by Stat5 of a factor that Stat1 requires to stimulate the IRF-1 promoter.

In addition to Stat5, Stat1 and Stat3 both have been reported to be activated by the PRLR (436, 501, 508). Stat3, also named APRF (acute phase response factor), was cloned as an IL-6-activated transcription factor (552). Stat1 was first isolated as part of the ISGF3 complex (containing Stat2 and a DNA-binding protein called p48) which is typically activated by IFNs α and β (553). Stat1 homodimers can also be formed upon IFN γ activation. The region(s) of the PRLR required for activation of these Stats remain poorly documented, although the 93 membrane-proximal residues have been reported to be sufficient for tyrosine phosphorylation of both Stats 1 and 3 (508). In the context of the GHR, it has recently been hypothesized that phosphotyrosine(s) of JAK2 could also bind to Stat3, in agreement with the presence of the consensus Stat3-binding site (see below) in the kinase (516, 520, 554, 555). Although such an interaction does not preclude the possible occurrence of interactions also with the receptor, this hypothesis remains to be tested in the context of Stat activation by the PRLR.

Stat-binding phosphotyrosine motifs in the PRLR. As described above, within a given receptor cytoplasmic domain, not all tyrosines undergo phosphorylation upon ligand-mediated receptor activation. The molecular features directing one tyrosine, and not another, to become phosphorylated are still poorly understood. Obviously, one requirement is the accessibility of the residue by the kinase. A phosphotyrosine must be recognized by at least three types of signaling proteins: the tyrosine kinase, the tyrosine phosphatase that will down-regulate the signal, and the SH2- (or any other motif recognizing phosphotyrosine) containing protein(s) that will dock on the phosphotyrosine. It is usually accepted that some of these interactions are driven by the recognition of specific sequences surrounding the phosphotyrosine (556, 557). Attempts have thus been made to correlate the ability of cytokine receptors to bind a given subset of Stat proteins with the presence of consensus sequences in the near environment of their cytoplasmic phosphotyrosines (520, 558–561).

Regarding Stat5 recruitment by the long PRLR, no sequence similarity could be found around the three phosphorylated tyrosines involved (Tyr 580, Tyr 479, Tyr 473; see above and Fig. 2B). The D-x-Y motif, involving some ty-

rosines within the GHR (516, 520), EPOR (548, 562, 563), and IL-2R β (544, 564), has been proposed as a putative Stat5-binding motif (520); none of the three phosphorylated tyrosines of the PRLR (514) matches such a consensus sequence. Recently, it has been suggested that an Asp at position -2 or -1, and a hydrophobic residue (preferentially a Leu or any aliphatic side chain) at positions +1 and +3 with respect to the phosphotyrosine could favor Stat 5 binding (561). Interestingly, two of the three proposed Stat5-binding sites in PRLR (Tyr 580 and 479) have an Asp at position -1, and all (including Tyr 473) have a hydrophobic residue at position +1 (Leu or Val). In contrast, the three other phosphotyrosines, which do not lead to Stat 5 activation, *i.e.*, Tyr 309, 402, and 515 (514) are in less agreement with this proposed consensus (Fig. 2B). In fact, the sequence bordering Tyr 402 is more closely related to that described for insulin receptor substrate-1 (IRS-1) binding.

Finally, we note the sequence bordering Tyr 309, which is phosphorylated in the context of a single-tyrosine PRLR mutant (514), matches the Y-x-x-Q sequence identified as Stat3-binding motif (558) and closely resembles that described for Stat1 binding (Y-P-x-Q instead of Y-x-P-Q; Fig. 2B) (559). This residue could thus be involved in Stat3 and/or Stat1 recruitment, in agreement with data obtained using a PRLR truncated at residue 328 (508). A similar proposal has been formulated recently by Yu-Lee and colleagues (482, 507) in view of the decrease of Stat1-mediated PRL induction of IRF-1 promoter activity. As an alternative to Stat/receptor direct interactions, such a consensus sequence is also present in JAK2 (555, 565).

Stat DNA-binding motifs. Consensus DNA motifs specifically recognized by Stat complexes have been identified in the promoters of target genes. The motif termed GAS (for γ -IFN-activated sequence) was defined using Stat homodimers and consists of a palindromic sequence TTC xxx GAA (532, 566). The specificity of the interaction between a particular Stat and a GAS motif found in a given target promoter has been proposed to depend, at least in part, on the center core nucleotide(s) (532). The activation of identical Stat proteins by different cytokine receptors questions the mechanisms by which specificity of signaling pathways is achieved in response to a particular hormonal stimulation. Although several cytokines (EPO, GM-CSF, GH, PRL, IL-2, IL-3, IL-5) activate the DNA-binding ability of Stat5 and/or transactivate the β -casein luciferase reporter gene *in vitro* (516, 536, 543, 564, 567-571), it is unlikely that all these cytokines stimulate the synthesis of this milk protein *in vivo*. This suggests that different Stat combinations and/or involvement of other signal transducers direct the specificity of the final response. For example, it has recently been reported that Stat5 interacts with the glucocorticoid receptor (572). This functional cooperation seems to require specific DNA binding of Stat5, but not of the glucocorticoid receptor (572a), although this hypothesis remains controversial (573).

In conclusion, particular emphasis must be given to the activation of Stat5, which is probably the major axis of the JAK-Stat cascade, if not of all signaling pathways activated by the PRLR, as confirmed by the very similar phenotypes analyzed thus far in PRLR, Stat5a and Stat5b knockout mice (see Section VII).

B. The Ras/Raf/MAP kinase pathway

Although the JAK-Stat cascade is presumably the most important signaling pathway used by cytokine receptors, other transducing pathways are also likely involved in signal transduction by these receptors. Signaling through MAP kinases (MAPK) involves the Shc/SOS/Grb2/Ras/Raf/MAPK cascade (574). Activation of the MAPK pathway has been reported in different cellular systems under PRL stimulation (157, 208, 443-445, 447, 525, 575). Whether activation of the MAP cascade requires JAK2, Fyn (or any Src kinase; see below), or any other pathway is currently unknown. Activation of the nucleotide exchange protein Vav has also been reported (446). Although the JAK-Stat and the MAPK cascades were initially regarded as independent pathways, several recent data suggest rather that they are interconnected (576).

C. Other signaling pathways

Fyn, a member of the Src kinase family (577), is associated with the PRLR and is activated by PRL stimulation in the rat T lymphoma Nb2 cell line (440). Association of the PRLR with Src, the prototype member of this kinase family, has also been reported after PRL stimulation in lactating rat hepatocytes (162). The role of Src kinases in signal transduction by PRLRs remains unknown, although promotion of cell growth has been suggested (162).

Using a modified rat PRLR cDNA encoding an additional N-terminal epitope specifically designed to allow the rapid purification of the receptor and associated proteins from transfected cells, we have recently shown that PRL induces a rapid tyrosine phosphorylation of IRS-1 and of the 85-kDa subunit of the phosphatidylinositol (PI)-3' kinase (450, 455). Both PI-3' kinase and IRS-1 appear to associate with the PRLR in a PRL-dependent manner. Recently, it has been proposed that PRL activation of PI-3' kinase might be mediated by Fyn in Nb2 cells (577a).

Since most transducer molecules are activated by tyrosine phosphorylation (JAKs, Stats, Src, etc.), involvement of tyrosine phosphatases to modulate or down-regulate the signaling cascades is expected. Accordingly, several recent studies pointed out a role of tyrosine phosphatases in PRLR signaling (451, 578, 579), although the mechanism by which they are regulated, as well as their substrates, remain poorly documented. We have reported that the phosphatase PTP-1D (now renamed SHP-2; see Ref. 580), is activated by JAK2 and acts as a positive regulator of PRLR-dependent induction of β -casein gene transcription (451). Moreover, a new family of SH2-containing protein inhibiting the JAK/Stat pathways has been recently identified (580a-580d). Members of this new protein family were named CIS (cytokine-inducible SH2-containing protein), SOCS (suppressor of cytokine signaling), JAB (JAK binding protein) and SSI (Stat-induced Stat inhibitor), and they inhibit the JAK/Stat pathway by interacting with JAKs (SOCS) or by competing with Stats for binding to the receptor (CIS). These proteins appear as targets of activated Stats and hence, provide a way to down-regulate the JAK/Stat pathways. PIAS3 (protein inhibitor of activated Stat) directly interacts with Stat3 and blocks its

binding to DNA targets (580e). Although involvement of these negative regulators of cytokine-activated JAK/Stat pathways has not been reported in connection with the PRLR, it is likely that at least some of these proteins are involved in negative feedback control of PRLR signaling.

Involvement of phospholipase-C (PLC γ) and PKC has been suggested, although the role of these enzymes in PRLR signaling remains unknown and their substrates remain poorly identified (155, 176, 177, 581). Finally, PRL has been shown to increase the concentration of intracellular calcium in PRLR-transfected CHO cells, although the physiological meaning of this phenomenon remains to be elucidated (582).

Cross-linking experiments have suggested that the PRLR is complexed to G proteins in Nb2 cells (583). The same group proposed the involvement of these guanine nucleotide-binding proteins in the mitogenic action of PRL on Nb2 cells (584). A potentially new signal transducer has been recently identified in rat ovary and appears to be specific to this tissue (585). This protein, named PRAP (PRLR-associated protein), preferentially binds to the short PRLR isoform and has been proposed to contribute to the luteotropic effects of PRL in the corpus luteum.

Finally, cross-talk between PRLR and EGF receptor (EGFR) has been recently suggested (585a, 585b). PRL could antagonize EGF signaling by increasing EGFR threonine phosphorylation and thereby decreasing EGF-induced EGFR tyrosine phosphorylation.

VII. Null Mutation of the PRLR Gene

Almost all of the effects attributed to PRL in vertebrates (92, 586) are the subject of conflicting observations and despite an amazing accumulation of articles, the precise role of lactogenic hormones, with the clear exception of direct mammary effects, remains an open question. Lactogens thus remain well characterized hormones with multiple but, in many cases, not well characterized functions.

In vivo, two animal models have been developed to study PRL action: one utilizes lowering PRL levels, achieved either by hypophysectomy or administration of dopamine D₂ receptor agonists; the second is based on the use of spontaneous mutant dwarf mice strains, lacking pituitary somatotrophs and lactotrophs. These models are, however, severely compromised by incomplete PRL depletion (extrapituitary PRL is not affected in these models; see Section II.D), and by the unavoidable suppression of other pituitary and nonpituitary hormones. *In vitro* models suffer similar limitations. FCS, an essential component of most cell and tissue culture media, contains high concentrations of lactogenic hormones that are difficult to remove (207). In addition, a growing list of cells (see Section II.D) synthesize and secrete PRL (27, 587, 588), making complete PRL removal impossible.

By the technique of gene targeting in mice (589), we have produced the first experimental model in which the effects of a complete absence of PRLR-mediated signaling of lactogenic hormones can be observed (214). Cell lines derived from this particular mouse offer the potential of *in vitro* models to further explore the indirect and direct effects of

PRL. The vast majority of mutations created by homologous recombination in embryonic stem cells to date have been null alleles. The next level of sophistication requires that a gene be inactivated or modified in specific tissues or at a certain time during the life of the animal (conditional knockouts or knockins).

The PRLR is expressed as short and long forms, differing in the length and sequence of their cytoplasmic tails (Section III.B; see Refs. 33, 46, 48, 50, and 51). The short and long forms are differentially expressed or regulated during the estrous cycle and pregnancy (52, 590, 591), which suggests that they may initiate distinct signaling pathways.

A. Gene cloning, vector construction, and generation of PRLR^{-/-} mice

A 129Sv mouse genomic DNA library was screened using the mouse PRLRS3 cDNA (51), and the coding region of the PRLR gene was isolated. Loss of just one cysteine of exons 4 or 5 encoding extracellular subdomain D1 results in complete lack of hormone-binding activity (66). A targeting construct was prepared with 7.5 kb of overall homology in which a 1.5-kb fragment containing exon 5 was replaced with the similarly sized Tk-NEO cassette, which resulted in a mutation creating an in-frame stop codon. After electroporation into E14.1 embryonic stem cells and neomycin selection, two selected clones were microinjected into 3.5-day-old C57BL/6 blastocysts and were able to generate germline chimeras. F1 intercrosses revealed a genotype distribution not significantly different from the normal Mendelian ratios excluding the first generation (214).

B. PRLR gene expression and PRLR protein in PRLR^{-/-} mice

By Northern blotting, a single major mRNA transcript of 2.8 kb was observed in PRLR^{+/+} but not in PRLR^{-/-} animals, demonstrating that no PRLR mRNA containing exon 5 is transcribed: the deletion of exon 5 was also confirmed by RT-PCR.

The PRLR protein was immunoprecipitated from solubilized liver microsomes and analyzed by immunoblot. A strong signal for the PRLR was detected using an antirat PRLR U5 (592) in the liver of PRLR^{+/+} mice; however, no PRLR protein could be detected in the liver of PRLR^{-/-} animals. PRLR^{-/-} mice showed no significant specific binding of PRL in liver microsomes. All data indicate that the exon 5 deletion caused the complete absence of functional PRLR in PRLR^{-/-} animals (214).

C. Impaired mammary gland development and lactation in heterozygous females

Most of the first litter of 6- to 8-week-old PRLR heterozygote (PRLR^{+/-}) F1 females died within 24 h, and the entire litter had perished by 48 h. The examination of the stomach contents of the pups showed air bubbles but no milk present, indicating that PRLR^{+/-} females were unable to lactate. This phenotype was not apparent after the second pregnancy, where all F1 PRLR^{+/-} females produced surviving pups. A similar phenotype was also seen at the first lactation in het-




FIG. 4. Histology of second thoracic mammary glands from $PRLR^{+/+}$ and $PRLR^{+/-}$ females. Hematoxylin/eosin-stained sections through the skin (left-hand side of each picture), subcutaneous fat (light staining), mammary fat pad (light staining), and epithelium (dark staining) of the number 2 and 3 glands (magnification, 100 \times). Alveoli are

eryozygous F2 females whatever the genetic backgrounds (C57BL/6 \times 129 and 129 F1). However, the 20-week-old $PRLR^{+/-}$ sisters of these animals produced their first litter with all surviving pups.

Histological examination of the mammary glands showed that lactational performance was correlated with the degree of mammary gland development (Fig. 4). These results demonstrate that two functional alleles of the $PRLR$ are required for efficient lactation and that this phenotype in heterozygotes is primarily due to a deficit in the degree of mammary gland development.

The mammary gland undergoes development *in utero*, at puberty (mainly ductal development), and during pregnancy (ductal and alveolar development). The essential hormonal factors regulating the later two phases in mice have been established to be estrogen, adrenocorticoids, and GH during puberty, and estrogen, progesterone, and PL and/or PRL during pregnancy (284, 593). These hormones produce some development with each estrous cycle and massive development at pregnancy, which never fully regresses after estrus or weaning, resulting in ever increasing alveolar and ductal development with each episode (594). Our observations suggest that the epithelial cell proliferation during pregnancy and the postpartum period depends on a threshold of $PRLR$ expression which is not achieved with just one functional gene, given that the level of $PRLR$ is closely controlled in mammary gland (595). In heterozygous mice where the level of the receptor is reduced, mammary gland proliferation is insufficient to ensure lactation at the first pregnancy but further estrous cycles or a single pregnancy lead to the development of a mammary gland capable of producing milk.

Initial histological investigation of the virgin gland of mature $PRLR^{-/-}$ animals indicated no dramatic differences as a result of the null mutation of the $PRLR$, with ductal tissue clearly present, confirming that the $PRLR$ is not essential for this stage of development similar to what has been reported for the null mutation of the progesterone receptor (596). The effect of this mutation on mammary development during pregnancy will be analyzed by transplantation of $PRLR^{-/-}$ mammary epithelium to $PRLR^{+/+}$ mammary fat pads cleared of endogenous epithelial cells before puberty.

D. Heterozygote maternal behavior

Some $PRLR^{+/-}$ mothers of 6–8 weeks and 20 weeks of age were observed to scatter their pups throughout the cage, often burying them completely in sawdust. When the mother reformed the nest, a pup was often left outside and not retrieved, while the others were suckled. This behavior was never observed among $PRLR^{+/+}$ and multiparous $PRLR^{+/-}$ females. In addition, $PRLR^{+/-}$ females generally did not eat their dead pups, in contrast to $PRLR^{+/+}$ females.

Other processes regulated by the hypothalamus, such as

indicated by *white arrows*, while ducts are indicated by *black arrows*. A, F1 $PRLR^{+/+}$ female 48 h post partum. Note engorged alveoli. B, F1 $PRLR^{+/-}$ female unable to lactate 48 h post partum. Note absence of alveoli and dominant ductal tissue. C, F1 $PRLR^{+/-}$ female showing partial lactation at 48 h post partum, with some functional alveoli present.




FIG. 5. Schematic representation of preimplantation egg development in $PRLR^{+/+}$ and $PRLR^{-/-}$ mice. On the left, normal egg development is represented. Undeveloped oocytes are depicted in white (including oocytes at the germinal vesicle stage, oocytes that have expelled the first polar body, and degenerated embryos), and 1 cell and 2 cell embryos, morula and blastocysts are color-coded. On the right, pie charts of the various stages of egg development at different times after ovulation are represented in $+/+$ vs. $-/-$ mice.

eating, sexual behavior, and locomotor activity, appear to be normal in $PRLR$ mutant mice.

E. Homozygous females are sterile

The $PRLR^{-/-}$ females mated irregularly, every 3 to 4 days, and never became pregnant. They show a number of reproductive deficiencies: all $PRLR^{-/-}$ females were sterile despite mating, which did not produce a pseudopregnancy. This was confirmed by examination of estrogen levels which showed a marked increase on day 3 after the vaginal plug, as the animals again entered estrus. The irregular mating patterns of the females indicate an alteration of estrous cyclicity. After mating, the PRL surges initiated by mechanical stimulation of the cervix are thought to induce pseudopregnancy (597), and the observation that the $PRLR^{-/-}$ females remated every 3 to 4 days demonstrates that a functional $PRLR$ is essential for the establishment of pseudopregnancy.

Multiple abnormalities were observed: fewer eggs were fertilized, oocytes at the germinal vesicle stage were released from the ovary, and fragmented embryos were found. The number of eggs ovulated was reduced, and histological investigation showed fewer primary follicles in $PRLR^{-/-}$ ovaries. Only 19% of blastocysts were recovered at day 3.5 in the uterus of $PRLR^{-/-}$ against 85% in wild-type animals (Fig. 5). Single cell-fertilized eggs were recovered, suggesting for most oocytes that an arrest of development occurred immediately after fertilization.

The $PRLR^{-/-}$ females were mated to $PRLR^{+/+}$ males, and fertilized embryos were reimplanted into the oviducts of pseudopregnant foster mothers. Most of the fertilized eggs

produced normal embryos. Using fertile $PRLR^{-/-}$ males (to exclude the possible paternal contribution of $PRLR$ to the embryo), embryos were also recovered and reached adulthood, demonstrating that the eggs are viable, and thus that lack of blastocyst implantation in $PRLR^{-/-}$ females may be due to a deficient environment in the oviduct.

Thus, the absence of $PRLR$ in female mice results in reduced ovulation, reduced fertilization, and almost complete arrest of preimplantation development. The blastocysts were unable to implant in $PRLR^{-/-}$ females, indicating that the uterus of these animals is refractory to implantation. The outcome is complete sterility.

The high number of eggs still containing germinal vesicles that were ovulated in $PRLR^{-/-}$ animals indicates that the $PRLR$ is important for oocyte maturation. Most of these were found on days 2.5 and 3.5 after the vaginal plug, suggesting that they may have been ovulated at a later time than those released at estrus; thus PRL may be involved in follicular atresia. A higher level of PRL is seen in follicles containing mature oocytes capable of being fertilized (337, 598–604), although others found no relationship (605) or suggest a negative effect (606). PRL was found to increase the rate of germinal vesicle breakdown and subsequent fertilization and correct development *in vitro* (607). The failure of a significant proportion of eggs to undergo germinal vesicle breakdown within maturing $PRLR^{-/-}$ follicles directly demonstrates the important influence of PRL and its receptor on oocyte maturation or atresia.

Fertilization rates were reduced, indicating that this effect is a result of a maternal deficiency. This may involve incom-

TABLE 8. Summary of phenotypes of PRLR^{+/-} and PRLR^{-/-} mice

	+/-	-/-
Reproduction		
Female	Normal mating, normal pregnancy	Sterility: No pseudopregnancy Ovulation rate decreased Oocyte maturation decreased No implantation Delayed fertility
Male	Normal	
Mammary gland development	Impaired development efficient lactation after the first pregnancy	No lactation: Ductal branching greatly reduced Absence of alveoli
Behavior		
Maternal	Reduced	Absent
Configural learning	Normal	Normal
Eating, locomotor activity, olfactory function	Normal	Normal
Immunity	Normal	No obvious differences

plete oocyte maturation, or a defect in the oviduct such as a reduced sperm transport to the ampulla or loss of a factor that enhances fertilization. Divergent effects of PRL on the rate of implantation development of mouse embryos have been reported (607–609). The present studies exclude the absolute requirement for an oocyte PRLR in pre- and postimplantation embryonic development, supporting previous investigations (610, 611) and indicating that the defect must reside in the environment in which the embryo develops. A number of factors in the oviduct that influence preimplantation development may be affected by the PRLR mutation. Estrogen and progesterone can influence the rate of ovum transport and preimplantation development (612).

The failure of trophic support of the corpus luteum by PRL would be expected to reduce progesterone levels on day 2.5 of pregnancy, when progesterone levels normally rise (613). Injection of antibodies against progesterone blocks mouse embryo development at the four-cell stage (614), but the failure of preimplantation development in PRLR^{-/-} females occurs earlier, when progesterone levels are normal in PRLR^{-/-} females. These results indicate that the PRLR must trigger a signal that occurs earlier than PRL-induced trophic support of the corpus luteum. Candidates include ovum factor, now identified as platelet activating factor (PAF), released by the fertilized eggs (615), and early pregnancy factor (EPF), a multifactorial activity comprised of PAF, thiodoxin (616), chaperonin 10 (617), and other uncharacterized molecules produced by the platelets of the oviduct and ovary in response to embryonic PAF (618). EPF is present in serum 24 h after ovulation and stimulates lymphocytes to produce a suppressor of the delayed-type hypersensitivity reaction, potentially protecting the ovum from the maternal immune system and promoting embryo cleavage, in addition to acting as a growth factor (619). Synthesis of EPF by isolated mouse oviducts and ovaries is stimulated by PRL, and PRL cooperates with PAF to stimulate ovarian EPF *in vivo* in response to fertilization, while passive immunization with antibodies against EPF produces almost identical effects as the PRLR mutation, with 54% of eggs not developing beyond the fertilized egg and two-cell stage (620–622).

Uterine preparation for embryo implantation is dependent upon continued estrogen and progesterone secretion by the corpus luteum of the ovary, which is supported by the pituitary in rodents during the first half of pregnancy (623). PRL has been shown to stimulate progesterone synthesis by dispersed ovarian cells from midpregnant mice (624), demonstrating that lactogenic hormones can directly stimulate ovarian progesterone secretion. Furthermore a nidatory ovarian estrogen surge is required to allow embryo implantation (625). Thus PRLR^{-/-} females probably cannot support the implantation of blastocysts because the corpus luteum does not receive pituitary PRL support, and thus progesterone- and estrogen-dependent signals for implantation cannot occur. Recently, multiple reproductive failures similar to those observed in PRLR knockout mice have been reported in cyclooxygenase 2-deficient mice (625a).

F. Homozygous male fertility

The fertility of PRLR^{-/-} males was examined by housing each male separately with a 12- to 14-week old PRLR^{+/-} female of proven fertility. At estrus, all females showed vaginal plugs, indicating normal mating, intromission, and ejaculation, but 20% of all the tested males showed a delayed fertility. Testes and accessory organs were of normal size; their histological examination showed no obvious morphological or histological abnormalities: clearly defined germinal cell layers and spermatocytes were present in the seminiferous tubules.

PRL may regulate testosterone production by Leydig cells via modulation of the effects of LH and of the level of its receptor (35). PRL has been also proposed to be involved in sperm capacitation (626): a short period of incubation with PRL has been reported to enhance *in vitro* fertilization rates (369), while longer periods reduce *in vitro* fertilization rates (610, 627), although others have seen no effect (611). PRL can also influence the function of the accessory reproductive glands (391, 628). The fact that if one waits long enough, all PRLR^{-/-} males are fully fertile, indicates that this role of PRL can be performed by other regulatory factors.

G. Other gene-targeted mutations leading to impaired mammary gland and reproductive functions

Although no effect on the mammary gland of heterozygous animals has been reported for null mutations of the progesterone receptor gene (596), the complete absence of this receptor in homozygous animals results in a gland lacking terminal-end buds with some branched ducts (596). Although estradiol receptor knockout females are infertile, to our knowledge, no particular mammary gland phenotype has been reported (629).

In Stat5a-deficient mice, mammary lobuloalveolar outgrowth during pregnancy was curtailed, and females were unable to lactate after parturition because of a failure of terminal differentiation (630). Similarly, mammary gland development is also impaired in Stat5b^{-/-} females and, although milk protein genes are expressed, there is insufficient milk to feed pups (630a). Similar phenotypes have been recently reported in PRL knockout mice (631). Interestingly, Stat5b, but not Stat5a-deficient females exhibit severely compromised fertility. Moreover, the phenotype of Stat5b knockout mice is distinct from that of Stat5a-deficient mice by a decrease in body growth profile in the former.

Mice homozygous for a germline mutation in *A-myb*, a nuclear protein regulator of transcription, show a marked underdevelopment of the breast epithelial compartment after pregnancy. Consequently, mice are unable to nurse their newborn pups, demonstrating a critical role of *A-myb* in mammary gland development (632). Mice lacking the cyclin D1 gene also exhibit a dramatic impairment of mammary gland development leading to inability to lactate their litters (633).

In conclusion, the phenotypes of animals lacking functional genes encoding PRLR, PRL, Stat5a, or Stat5b confirm the essential role of these molecules in the signaling pathway(s) leading to mammary gland development and/or reproductive function, whereas estradiol and progesterone receptors are probably involved to a lesser extent in mammary gland development.

H. Other phenotypes of PRLR^{-/-} mice

Immunoregulation. In view of the putative immunomodulatory role of PRL, we are currently analyzing the immune phenotype of the knockout animals. Preliminary data suggest that maturation and export of precursor cells occur in the thymus and, to date, no abnormality in the export of the lymphoid system to the periphery has been identified. Interestingly, no immune phenotype was seen in mice lacking the PRL gene (631). Specific functional studies to characterize individual immune cell types in PRLR^{-/-} are ongoing.

Bone. The PRLR mRNA expression has been detected in murine bone cells. The level of the mRNA encoding the long form of PRLR in osteoblasts is comparable to that observed in other cells such as thymocytes, mammary cells, or bone marrow cells. No expression of any form was found in osteoclast-like cells. In initial studies, examination of the calvariae of PRLR^{-/-} embryos at 18.5 days of age indicates that these bones are less developed or more disorganized than in wild-type controls.

Late fetal or neonatal lethality was the expected phenotype of PRLR^{-/-} animals. PRLR expression rises dramatically in a number of rodent tissues during the late stages of pregnancy, and PRL and PLs are detectable in fetal blood (634, 635), suggesting that lactogenic hormones begin to exert major effects during this period in preparation for the transfer to autonomous life. In contrast, PRLR^{-/-} animals were born and survived until adulthood. A number of phenotypes were observed in both hetero- and homozygous animals (see Table 8). It is probable that most of the phenotypes observed are related to the absence of the long form of the receptor, since this is the major isoform in all cells involved in reproductive function. Almost every aspect of female reproduction is altered in these animals, unambiguously demonstrating that the PRLR is a key regulator of reproduction. The ability of this new model to provide novel insights into the function of lactogenic hormones and their receptor illustrates the power of the knockout approach to discover and confirm specific roles for well investigated molecules. The generation of knockouts has highlighted the role of many genes in embryonic development, yet the study of phenotypes associated with null alleles is only an initial step in the analysis of the gene function. We can look forward, in the near future, to the widespread application of approaches of greater technical sophistication, including the generation of subtle alterations in the gene sequence and conditional knockouts.

VIII. Summary and Conclusions

PRL is an anterior pituitary hormone that, along with GH and PLs, forms a family of hormones that probably resulted from the duplication of an ancestral gene. The PRLR is also a member of a larger family, known as the cytokine class-1 receptor superfamily, which currently has more than 20 different members. PRLRs or binding sites are widely distributed throughout the body. In fact, it is difficult to find a tissue that does not express any PRLR mRNA or protein. In agreement with this wide distribution of receptors is the fact that now more than 300 separate actions of PRL have been reported in various vertebrates, including effects on water and salt balance, growth and development, endocrinology and metabolism, brain and behavior, reproduction, and immune regulation and protection. Clearly, a large proportion of these actions are directly or indirectly associated with the process of reproduction, including many behavioral effects. PRL is also becoming well known as an important regulator of immune function. A number of disease states, including the growth of different forms of cancer as well as various autoimmune diseases, appear to be related to an overproduction of PRL, which may act in an endocrine, autocrine, or paracrine manner, or via an increased sensitivity to the hormone.

The first step in the mechanism of action of PRL is the binding to a cell surface receptor. The ligand binds in a two-step process in which site 1 on PRL binds to one receptor molecule, after which a second receptor molecule binds to site 2 on the hormone, forming a homodimer consisting of one molecule of PRL and two molecules of receptor. The PRLR contains no intrinsic tyrosine kinase cytoplasmic do-

main but associates with a cytoplasmic tyrosine kinase, JAK2. Dimerization of the receptor induces tyrosine phosphorylation and activation of the JAK kinase followed by phosphorylation of the receptor. Other receptor-associated kinases of the Src family have also been shown to be activated by PRL. One major pathway of signaling involves phosphorylation of cytoplasmic Stat proteins, which themselves dimerize and translocate to nucleus and bind to specific promoter elements on PRL-responsive genes. In addition, the Ras/Raf/MAP kinase pathway is also activated by PRL and may be involved in the proliferative effects of the hormone. Finally, a number of other potential mediators have been identified, including IRS-1, PI-3 kinase, SHP-2, PLC γ , PKC, and intracellular Ca²⁺.

The technique of gene targeting in mice has been used to develop the first experimental model in which the effects of the complete absence of any lactogen or PRL-mediated effects can be studied. Heterozygous (+/-) females show almost complete failure to lactate after the first, but not subsequent, pregnancies. Homozygous (-/-) females are infertile due to multiple reproductive abnormalities, including ovulation of premeiotic oocytes, reduced fertilization of oocytes, reduced preimplantation oocyte development, lack of embryo implantation, and the absence of pseudopregnancy. Twenty per cent of the homozygous males showed delayed fertility. Other phenotypes, including effects on the immune system and bone, are currently being examined.

It is clear that there are multiple actions associated with PRL. It will be important to correlate known effects with local production of PRL to differentiate classic endocrine from autocrine/paracrine effects. The fact that extrapituitary PRL can, under some circumstances, compensate for pituitary PRL raises the interesting possibility that there may be effects of PRL other than those originally observed in hypophysectomized rats. The PRLR knockout mouse model should be an interesting system by which to look for effects activated only by PRL or other lactogenic hormones. On the other hand, many of the effects reported in this review may be shared with other hormones, cytokines, or growth factors and thus will be more difficult to study.

Although PRL evolved several hundred million years ago, we are now at the end of the 20th century just beginning to understand how the hormone acts and its potential involvement in pathological disease states. Future research will center on further expanding the already long list of PRL actions and attempt to better understand the mechanisms of action of this intriguing hormone.

Acknowledgments

The authors are grateful to Claudine Coridun for help in the preparation of this manuscript and to Alain Pezet, Philippe Clément-Lacoix, and Sandrina Kinet for preparation of the figures. We thank Drs. P. Elkins and A.M. de Vos for providing the figure of the ECD of the PRLR.

References

1. Stricker P, Grueter R 1928 Action du lobe antérieur de l'hypophyse sur la montée laiteuse. *C R Soc Biol* 99:1978-1980
2. Riddle O, Bates RW, Dykshorn SW 1933 The preparation, identification and assay of prolactin - a hormone of the anterior pituitary. *Am J Physiol* 105:191-216
3. Hwang P, Guyda H, Friesen H 1971 A radioimmunoassay for human prolactin. *Proc Natl Acad Sci USA* 68:1902-1906
4. Lewis UJ, Singh RN, Seavey BK 1971 Human prolactin: isolation and some properties. *Biochem Biophys Res Commun* 44:1169-1176
5. Li CH, Dixon JS, Lo TB, Pankov YM, Schmidt KD 1969 Amino acid sequence of ovine lactogenic hormone. *Nature* 224:695-696
6. Nicoll CS, Mayer GL, Russell SM 1986 Structural features of prolactins and growth hormones that can be related to their biological properties. *Endocr Rev* 7:169-203
7. Niall HD, Hogan ML, Sauer R, Rosenblum IV, Greenwood FC 1971 Sequences of pituitary and placental lactogenic and growth hormones: evolution from a primordial peptide by gene duplication. *Proc Natl Acad Sci USA* 68:866-870
8. Miller WL, Eberhardt NL 1983 Structure and evolution of the growth hormone gene family. *Endocr Rev* 4:97-130
9. Goffin V, Shiverick KT, Kelly PA, Martial JA 1996 Sequence-function relationships within the expanding family of prolactin, growth hormone, placental lactogen and related proteins in mammals. *Endocr Rev* 17:385-410
10. Owerbach D, Rutter WJ, Cooke NE, Martial JA, Shows TB 1981 The prolactin gene is located on chromosome 6 in humans. *Science* 212:815-816
11. Truong AT, Duez C, Belayew A, Renard A, Pictet R, Bell GI, Martial JA 1984 Isolation and characterization of the human prolactin gene. *EMBO J* 3:429-437
12. Cooke NE, Coit D, Shine J, Baxter JD, Martial JA 1981 Human prolactin: cDNA structural analysis and evolutionary comparisons. *J Biol Chem* 256:4007-4016
13. Cooke NE, Coit D, Weiner RI, Baxter JD, Martial JA 1980 Structure of cloned DNA complementary to rat prolactin messenger RNA. *J Biol Chem* 255:6502-6510
14. Miller WL, Coit D, Baxter JD, Martial JA 1981 Cloning of bovine prolactin cDNA and evolutionary implications of its sequence. *DNA* 1:37-50
15. Mercier L, Rentier-Delrue F, Swennen D, Lion M, Le Goff P, Prunet P, Martial JA 1989 Rainbow trout prolactin cDNA cloning in *Escherichia coli*. *DNA* 8:119-125
16. Rentier-Delrue F, Swennen D, Prunet P, Lion M, Martial JA 1989 Tilapia prolactin: molecular cloning of two cDNAs and expression in *Escherichia coli*. *DNA* 8:261-270
17. Sinha YN 1995 Structural variants of prolactin: occurrence and physiological significance. *Endocr Rev* 16:354-369
18. Yasuda A, Miyazima KI, Kawauchi H, Peter RE, Lin HR, Yamaguchi K, Sano H 1987 Primary structure of common carp prolactins. *Gen Comp Endocrinol* 66:280-290
19. Walker AM 1994 Phosphorylated and non-phosphorylated prolactin isoforms. *Trends Endocrinol Metab* 5:195-200
20. Martial JA, Cooke NE 1980 Isolation and sequence analysis of the rat prolactin structural gene. In: McLeod RM, Scapagnini U (eds) *Central and Peripheral Regulation of Prolactin Function*. Raven Press, New York, pp 115-123
21. Rand-Weaver M, Kawauchi H 1993 Growth hormone, prolactin and somatolactin: a structural overview. In: Hochachka PW, Mammens TP (eds) *Biochemistry and Molecular Biology of Fishes*. Elsevier Science Publishers, Amsterdam, pp 39-56
22. Walker WH, Fitzpatrick SL, Barrera-Saldana HA, Resendez-Perez D, Saunders GF 1991 The human placental lactogen genes: structure, function, evolution and transcriptional regulation. *Endocr Rev* 12:316-328
23. Bewley TA, Li CH 1972 Circular dichroism studies on human pituitary growth hormone and ovine pituitary lactogenic hormone. *Biochemistry* 11:884-888
24. Goffin V, Martial JA, Summers NL 1995 Use of a model to understand prolactin and growth hormone specificities. *Protein Eng* 8:1215-1231
25. Abdel-Meguid SS, Shieh HS, Smith WW, Dayringer HE, Violand BN, Bentle LA 1987 Three-dimensional structure of a genetically engineered variant of porcine growth hormone. *Proc Natl Acad Sci USA* 84:6434-6437
26. De Vos AM, Ultsch M, Kossiakoff AA 1992 Human growth hor-

- mon and extracellular domain of its receptor: crystal structure of the complex. *Science* 255:306–312
27. Ben-Jonathan N, Mershon JL, Allen DL, Steinmetz RW 1996 Extrahypothalamic prolactin: distribution, regulation, functions, and clinical aspects. *Endocr Rev* 17:639–669
 28. Nagy E, Berczi I 1991 Hypophysectomized rats depend on residual prolactin for survival. *Endocrinology* 128:2776–2784
 29. Posner BI, Kelly PA, Shiu RP, Friesen HG 1974 Studies of insulin, growth hormone and prolactin binding: tissue distribution, species variation and characterization. *Endocrinology* 95:521–531
 30. Posner BI, Kelly PA, Friesen HG 1975 Prolactin receptors in rat liver: possible induction by prolactin. *Science* 188:57–59
 31. Kelly PA, Bradley C, Shiu RP, Meites J, Friesen HG 1974 Prolactin binding to rat mammary tumor tissue. *Proc Soc Exp Biol Med* 146:816–819
 32. Kelly PA, Posner BI, Tsushima T, Shiu RP, Friesen HG 1974 Tissue distribution and ontogeny of growth hormone and prolactin receptors. In: Raiti S (ed) *Advances in Human Growth Hormone Research*. General Printing Office, Washington, DC, pp 567–584
 33. Boutin JM, Jolicoeur C, Okamura H, Gagnon J, Edery M, Shirota M, Banville D, Dusanter-Fourt I, Djiane J, Kelly PA 1988 Cloning and expression of the rat prolactin receptor, a member of the growth hormone/prolactin receptor gene family. *Cell* 53:69–77
 34. Leung DW, Spencer SA, Cachianes G, Hammonds RG, Collins C, Henzel WJ, Barnard R, Waters MJ, Wood WI 1987 Growth hormone receptor and serum binding protein: purification, cloning, and expression. *Nature* 330:537–543
 35. Kelly PA, Djiane J, Postel-Vinay MC, Edery M 1991 The prolactin/growth hormone receptor family. *Endocr Rev* 12:235–251
 36. Goffin V, Kelly PA 1996 Prolactin and growth hormone receptors. *Clin Endocrinol (Oxf)* 45:247–255
 37. Goffin V, Kelly PA 1996 The prolactin/growth hormone receptor family: structure/function relationships. *J Mammary Gland Biol Neopl* 2:7–17
 38. Goffin V, Ferrag F, Kelly PA 1998 Molecular aspects of prolactin and growth hormone receptors. In: LeRoith D (ed) *Advances in Mol Cell Endocrinol*, Vol 2. JAI Press, pp 1–33
 39. Bazan F 1989 A novel family of growth factor receptors: a common binding domain in the growth hormone, prolactin, erythropoietin and IL-6 receptors, and p75 IL-2 receptor β -chain. *Biochem Biophys Res Commun* 164:788–795
 40. Bazan JF 1990 Structural design of molecular evolution of a cytokine receptor superfamily. *Proc Natl Acad Sci USA* 87:6934–6938
 41. Taga T, Kishimoto T 1992 Cytokine receptors and signal transduction. *FASEB J* 6:3387–3396
 42. Stahl N, Yancopoulos GD 1993 The alphas, betas and kinases of cytokine receptor complexes. *Cell* 74:587–590
 43. Kitamura T, Ogorochi T, Miyajima A 1994 Multimeric cytokine receptors. *Trends Endocrinol Metab* 5:8–14
 44. Finidori J, Kelly PA 1995 Cytokine receptor signalling through two novel families of transducer molecules: Janus kinases, and signal transducers and activators of transcription. *J Endocrinol* 147:11–23
 45. Wells JA, De Vos AM 1996 Hematopoietic receptor complexes. *Annu Rev Biochem* 65:609–634
 46. Arden KC, Boutin J-M, Djiane J, Kelly PA, Cavenne WK 1990 The receptors for prolactin and growth hormone are localized in the same region of human chromosome 5. *Cytogenet Cell Genet* 53:161–165
 47. Ali S, Pellegrini I, Edery M, Lesueur L, Paly J, Djiane J, Kelly PA 1991 A prolactin-dependent immune cell line (Nb2) expresses a mutant form of prolactin receptor. *J Biol Chem* 266:20110–20117
 48. Boutin JM, Edery M, Shirota M, Jolicoeur C, Lesueur L, Ali S, Gould D, Djiane J, Kelly PA 1989 Identification of a cDNA encoding a long form of prolactin receptor in human hepatoma and breast cancer cells. *Mol Endocrinol* 3:1455–1461
 49. Nagano M, Chastre E, Choquet A, Bara J, Gespach C, Kelly PA 1995 Expression of prolactin and growth hormone receptor genes and their isoforms in the gastrointestinal tract. *Am J Physiol* 268:G431–G442
 50. Shirota M, Banville D, Ali S, Jolicoeur C, Boutin J-M, Edery M, Djiane J, Kelly PA 1990 Expression of two forms of prolactin receptor in rat ovary and liver. *Mol Endocrinol* 4:1136–1143
 51. Davis JA, Linzer DIH 1989 Expression of multiple forms of the prolactin receptor. *Mol Endocrinol* 3:674–680
 52. Clarke DL, Linzer DIH 1993 Changes in prolactin receptor expression during pregnancy in the mouse ovary. *Endocrinology* 133:224–232
 53. Postel-Vinay MC, Belair L, Kayser C, Kelly PA, Djiane J 1991 Identification of prolactin and growth hormone binding proteins in milk. *Proc Natl Acad Sci USA* 88:6687–6690
 54. Fuh G, Wells JA 1995 Prolactin receptor antagonists that inhibit the growth of breast cancer cell lines. *J Biol Chem* 270:13133–13137
 55. Amit T, Dibner C, Barkey RJ 1997 Characterization of prolactin- and growth hormone-binding proteins in milk and their diversity among species. *Mol Cell Endocrinol* 130:167–180
 56. Postel-Vinay MC 1996 Growth hormone- and prolactin-binding proteins: soluble forms of receptors. *Horm Res* 45:178–181
 57. Kossiakoff AA, Somers W, Ultsch M, Andow K, Muller YA, De Vos AM 1994 Comparison of the intermediate complexes of human growth hormone bound to the human growth hormone and prolactin receptors. *Protein Sci* 3:1697–1705
 58. Chen X, Horseman ND 1994 Cloning, expression and mutational analysis of the pigeon prolactin receptor. *Endocrinology* 135:269–276
 59. Gao J, Hughes JP, Auferin B, Buteau H, Edery M, Zhuang H, Wojchowski DM, Horseman ND 1996 Interactions among Janus kinases and the prolactin (PRL) receptor in the regulation of a PRL response element. *Mol Endocrinol* 10:847–856
 60. Murakami M, Narazaki M, Hibi M, Yawata H, Yasukawa K, Hamaguchi M, Taga T, Kishimoto T 1991 Critical cytoplasmic region of the interleukin-6 signal transducer, gp 130, is conserved in the cytokine receptor family. *Proc Natl Acad Sci USA* 88:11349–11353
 61. Vincent V, Goffin V, Rozakis-Adcock M, Mornon JP, Kelly PA 1997 Identification of cytoplasmic motifs required for short prolactin receptor internalization. *J Biol Chem* 272:7062–7068
 62. Somers W, Ultsch M, De Vos AM, Kossiakoff AA 1994 The X-ray structure of the growth hormone-prolactin receptor complex. *Nature* 372:478–481
 63. Caravella JA, Lyne PD, Richards WG 1996 A partial model of the erythropoietin receptor complex. *Proteins* 24:394–401
 64. Livnah O, Stura EA, Johnson DL, Middleton SA, Mulcahy LS, Wrighton NC, Dower WJ, Jolliffe LK, Wilson IA 1996 Functional mimicry of a protein hormone by a peptide agonist: the EPO receptor complex at 2.8 Å. *Science* 273:464–471
 65. Walter MR, Windsor WT, Nagabhushan TL, Lundell DJ, Lunn CA, Zaudodny PJ, Narula SK 1995 Crystal structure of a complex between interferon- γ and its soluble high-affinity receptor. *Nature* 376:230–235
 66. Rozakis-Adcock M, Kelly PA 1991 Mutational analysis of the ligand binding domain of the prolactin receptor. *J Biol Chem* 266:16472–16477
 67. Rozakis-Adcock M, Kelly PA 1992 Identification of ligand binding determinants of the prolactin receptor. *J Biol Chem* 267:7428–7433
 68. Bass SH, Mulkerrin MG, Wells JA 1991 A systematic mutational analysis of hormone-binding determinants in the human growth hormone receptor. *Proc Natl Acad Sci USA* 88:4498–4502
 69. Baumgartner JW, Wells CA, Chen CM, Waters MJ 1994 The role of the WSxWS motif in growth hormone receptor function. *J Biol Chem* 269:29094–29101
 70. Miyazaki T, Maruyama M, Yamada G, Hatakeyama M, Taniguchi T 1991 The integrity of the conserved WS motif common to IL-2 and other cytokine receptors is essential for ligand binding and signal transduction. *EMBO J* 10:3191–3197
 71. Quelle DE, Quelle FW, Wojchowski DM 1992 Mutations in the WSAWSE and cytosolic domains of the erythropoietin receptor affect signal transduction and ligand binding and internalization. *Mol Cell Biol* 12:4553–4561
 72. Hilton DJ, Watowich SS, Katz L, Lodish HF 1996 Saturation mutagenesis of the WSXWS motif of the erythropoietin receptor. *J Biol Chem* 271:4699–4708
 73. Sandowski Y, Nagano M, Bignon C, Djiane J, Kelly PA, Gertler A 1995 Preparation and characterization of recombinant prolactin receptor extracellular domain from rat. *Mol Cell Endocrinol* 115:1–11

- 73a. Buteau H, Pezet A, Ferrag F, Perrot-Applanat M, Kelly PA, Ederly M 1998 N-glycosylation of the prolactin receptor is not required for activation of gene transcription but is critical for its cell surface targeting. *Mol Endocrinol* 12:544-555
74. Gertler A, Petridou B, Kriwi GG, Djiane J 1993 Interaction of lactogenic hormones with purified recombinant extracellular domain of rabbit prolactin receptor expressed in insect cells. *FEBS Lett* 319:277-281
75. Hooper KP, Padmanabhan R, Ebner KE 1993 Expression of the extracellular domain of the rat liver prolactin receptor and its interaction with ovine prolactin. *J Biol Chem* 268:22347-22352
76. Goffin V, Norman M, Martial JA 1992 Alanine-scanning mutagenesis of human prolactin: importance of the 58-74 region for bioactivity. *Mol Endocrinol* 6:1381-1392
77. Goffin V, Struman I, Goormaghtigh E, Martial JA 1993 The addition of nine residues at the C-terminus of human prolactin drastically alters its biological properties. *Eur J Biochem* 214:483-490
78. Kinet S, Goffin V, Mainfroid V, Martial JA 1996 Characterization of lactogen receptor binding site 1 of human prolactin. *J Biol Chem* 271:14353-14360
79. Goffin V, Struman I, Mainfroid V, Kinet S, Martial JA 1994 Evidence for a second receptor binding site on human prolactin. *J Biol Chem* 269:32598-32606
80. Goffin V, Kinet S, Ferrag F, Binart N, Martial JA, Kelly PA 1996 Antagonistic properties of human prolactin analogs that show paradoxical agonistic activity in the Nb2 bioassay. *J Biol Chem* 271:16573-16579
81. Gertler A, Grosclaude J, Strasburger CJ, Nir S, Djiane J 1996 Real-time kinetic measurements of the interactions between lactogenic hormones and prolactin-receptor extracellular domains from several species support the model of hormone-induced transient receptor dimerization. *J Biol Chem* 271:24482-24491
82. Ozenberger BA, Young KH 1995 Functional interaction of ligands and receptors of the hematopoietic superfamily in yeast. *Mol Endocrinol* 9:1321-1329
83. Wells JA 1994 Structural and functional basis for hormone binding and receptor oligomerization. *Curr Opin Cell Biol* 6:163-173
84. Wells JA 1996 Binding in the growth hormone receptor complex. *Proc Natl Acad Sci USA* 93:1-6
85. Nagano M, Kelly PA 1994 Tissue distribution and regulation of rat prolactin receptor gene expression: quantitative analysis by polymerase chain reaction. *J Biol Chem* 269:13337-13345
86. Ouhitit A, Morel G, Kelly PA 1993 Visualization of gene expression of short and long forms of prolactin receptor in the rat. *Endocrinology* 133:135-144
87. Ouhitit A, Morel G, Kelly PA 1993 Visualization of gene expression of short and long forms of prolactin receptor in rat reproductive tissues. *Biol Reprod* 49:528-536
88. Ouhitit A, Kelly PA, Morel G 1994 Visualization of gene expression of short and long forms of prolactin receptor in rat digestive tissues. *Am J Physiol* 266:G807-G815
89. Royster M, Driscoll P, Kelly PA, Freemark M 1995 The prolactin receptor in the fetal rat: cellular localization of messenger RNA, immunoreactive protein, and ligand binding activity and induction of expression in late gestation. *Endocrinology* 136:3892-3900
90. Freemark M, Driscoll P, Andrews J, Kelly PA, Royster M 1996 Ontogenesis of prolactin receptor gene expression in the rat olfactory system: potential roles for lactogenic hormones in olfactory development. *Endocrinology* 137:934-942
91. Astwood EB 1941 The regulation of corpus luteum function by hypophysial luteotrophin. *Endocrinology* 29:309-319
92. Nicoll CS, Bern H 1972 On the actions of PRL among the vertebrates: is there a common denominator? In: Wolstenholme GEW, Knight J (eds) *Lactogenic Hormones*. Churchill Livingstone, London, pp 299-337
93. Nicoll CS 1974 Physiological actions of prolactin. In: Knobil E, Sawyer WH (eds) *Handbook of Physiology*. Section 7: Endocrinology. American Physiological Society, Washington, DC, pp 253-292
94. Maetz J 1970 Mechanisms of salt and water transfer across membranes in teleosts in relation to the aquatic environment. *Mem Soc Endocrinol* 18:3-29
95. Ogawa M, Yagasaki M, Yamazaki F 1973 The effect of prolactin on water influx in isolated gills of the goldfish, *Carassius auratus* L. *Comp Biochem Physiol A* 44:1177-1183
96. Leatherland JF, Lam TJ 1969 Effect of prolactin on the density of mucous cells on the gill filaments of the marine form (*trachurus*) of the threespine stickleback, *Gasterosteus aculeatus* L. *Can J Zool* 47:787-792
97. Pickford GE, Griffith RW, Torreti J, Hendlez E, Epstein FH 1970 Branchial reduction and renal stimulation of (Na^+ , K^+)-ATPase by prolactin in hypophysectomized killifish in fresh water. *Nature* 228:378-379
98. Flik G, Fenwick JC, Wendelaar Bonga SE 1989 Calcitropic actions of prolactin in fresh water North American eel (*Anguilla rostrata* LeSueur). *Am J Physiol* 257:R74-R79
99. Flik G, Rentier-Delrue F, Wendelaar Bonga SE 1994 Calcitropic effects of recombinant prolactins in *Oreochromis mossambicus*. *Am J Physiol* 266:1302-1308
100. Lam TJ, Leatherland JF 1969 Effects of prolactin on the glomerulus of the marine threespine stickleback, *Gasterosteus aculeatus* L., form *trachurus*, after transfer from seawater to fresh water, during the late autumn and early winter. *Can J Zool* 47:245-250
101. Olivereau M, Lemoine AM 1968 Effect of prolactin on the intact eel. 3. Effect on the histological structure of the kidney. *Z Zellforsch Mikrosk Anat* 88:576-590
102. Pippard C, Baylis PH 1986 Prolactin stimulates Na^+ - K^+ -ATPase activity located in the outer renal medulla of the rat. *J Endocrinol* 108:95-99
103. Richardson BP 1973 Evidence for a physiological role of prolactin in osmoregulation in the rat after its inhibition by 2-bromo- α -ergokryptine. *Br J Pharmacol* 47:623P-624P
104. Brown SC, Horgan EA, Savage LM, Brown PS 1986 Changes in body water and plasma constituents during bullfrog development: effects of temperature and hormones. *J Exp Zool* 237:25-33
105. Roberts JR, Dantzer WH 1992 Micropuncture study of avian kidney: effect of prolactin. *Am J Physiol* 262:933-937
106. Hirano T, Johnson DW, Bern HA 1971 Control of water movement in flounder urinary bladder by prolactin. *Nature* 230:469-470
107. Snart RS, Dalton T 1973 Response of toad bladder to prolactin. *Comp Biochem Physiol* 45:307-311
108. Ogawa M 1970 Effects of prolactin on the epidermal mucous cells of the goldfish, *Carassius auratus* L. *Can J Zool* 48:501-503
109. Moriya T 1982 Prolactin induces increase in the specific gravity of salamander, *Hynobius retardatus*, that raises adaptability to water. *J Exp Zool* 223:83-88
110. Hasegawa S, Hirano T, Kawauchi H 1986 Sodium-retaining activity of chum salmon prolactin in some euryhaline teleosts. *Gen Comp Endocrinol* 63:309-317
111. Robertson MT, Boyajian MJ, Patterson K, Robertson WV 1986 Modulation of the chloride concentration of human sweat by prolactin. *Endocrinology* 119:2439-2444
112. Mainoya JR, Bern HA, Regan JW 1974 Influence of ovine prolactin on transport of fluid and sodium chloride by the mammalian intestine and gall bladder. *J Endocrinol* 63:311-317
113. Mainoya JR 1975 Effects of bovine growth hormone, human placental lactogen and ovine prolactin on intestinal fluid and ion transport in the rat. *Endocrinology* 96:1165-1170
114. Mainoya JR 1975 Further studies on the action of prolactin on fluid and ion absorption by the rat jejunum. *Endocrinology* 96:1158-1164
115. Mainoya JR 1975 Analysis of the role of endogenous prolactin on fluid and sodium chloride absorption by the rat jejunum. *J Endocrinol* 67:343-349
116. Mainoya JR 1979 Effect of prolactin on fluid and NaCl absorption by the rat proximal and distal colon. *Experientia* 35:1060-1061
117. Peaker MJ, Phillips JG, Wright A 1970 The effect of prolactin on the secretory activity of the nasal salt-gland of the domestic duck (*Anas platythynchos*). *J Endocrinol* 47:123-127
118. Young KH, Kraeling RR, Bazer FW 1989 Effects of prolactin on conceptus survival and uterine secretory activity in pigs. *J Reprod Fertil* 86:713-722
119. Josimovich JB, Merisko K, Boccella L 1977 Amniotic prolactin control over amniotic and fetal extracellular fluid water and electrolytes in the rhesus monkey. *Endocrinology* 100:564-570
120. Sarandakou A, Kassanos D, Phocas I, Kontoravdis A, Chryssicopoulos A, Zourlas PA 1992 Amniotic fluid hormone profiles

- during normal and abnormal pregnancy. Clin Exp Obstet Gynecol 19:180-188
121. Kulczycki LL, Robertson MT 1988 The sweat chloride concentration and prolactin activity in cystic fibrosis. Scand J Gastroenterol 143:28-30
 122. Silverin B 1980 Effects of prolactin on the gonad and body weight of the male pied flycatcher during the breeding period. Endokrinologie 76:45-50
 123. Pérez-Villamil B, Bordiu E, Puente-Cueva M 1992 Involvement of physiological prolactin levels in growth and prolactin receptor content of prostate glands and testes in developing male rats. J Endocrinol 132:449-459
 124. Rynikova A, Koppel J, Kuchar S, Cikos S, Mozes S 1988 Effects of ovine prolactin in infant rats. Exp Clin Endocrinol 92:241-244
 125. De Jesus EG, Hirano T, Inui Y 1994 The antimetamorphic effect of prolactin in the Japanese flounder. Gen Comp Endocrinol 93:44-50
 126. Dodd MHI, Dodd JM 1976 The biology of metamorphosis. In: Lofts B (ed) Physiology of the Amphibia. Academic Press, New York, pp 467-599
 127. Yoshizato K, Yasumasu I 1970 Effect of prolactin on the tadpole tail fin. I. Stimulatory effect of prolactin on the collagen synthesis of the tadpole tail fin. Dev Growth Differ 11:305-317
 128. Yamamoto K, Kikuyama S 1982 Effect on prolactin antiserum on growth and resorption of tadpole tail. Endocrinol Jpn 29:81-85
 129. Berman RH, Bern HA, Nicoll CS, Strohman RC 1964 Growth-promoting effects of mammalian prolactin and growth hormone in tadpoles of *Rana catesbeiana*. J Exp Zool 156:353-360
 130. Bres O, Nicoll CS 1993 Effects of prolactin, growth hormone and triiodothyronine on prolactin receptors in larval and adult tiger salamanders (*Ambystoma tigrinum*). J Exp Zool 266:290-298
 131. Wright ML, Cykowski LJ, Lundrigan L, Hemond K, Kochan DM, Faszewski EE, Anuszewski CM 1994 Anterior pituitary and adrenal cortical hormones accelerate or inhibit tadpole hindlimb growth and development depending on stage of spontaneous development or thyroxine concentration in induced metamorphosis. J Exp Zool 270:175-188
 132. Tata JR, Kawahara A, Baker BS 1991 Prolactin inhibits both thyroid hormone-induced morphogenesis and cell death in cultured amphibian larval tissues. Dev Biol 146:72-80
 133. Clemons GK, Nicoll CS 1977 Effects of antisera to bullfrog prolactin and growth hormone on metamorphosis of *Rana catesbeiana* tadpoles. Gen Comp Endocrinol 31:495-497
 134. Tata JR 1993 Gene expression during metamorphosis: an ideal model for post-embryonic development. Bioessays 15:239-248
 135. Dent JN, Eng LA, Forbes MS 1973 Relations of prolactin and thyroid hormone to molting, skin texture, and cutaneous secretion in the red-spotted newt. J Exp Zool 184:369-382
 136. Hutchison RE 1975 Effects of ovarian steroids and prolactin on the sequential development of nesting behaviour in female budgerigars. J Endocrinol 67:29-39
 137. Schwabl H, Schwabl-Benzinger I, Goldsmith AR, Farner DS 1988 Effects of ovariectomy on long-day-induced premigratory fat deposition, plasma levels of luteinizing hormone and prolactin, and molt in white-crowned sparrows, *Zonotrichia leucophrys gambelii*. Gen Comp Endocrinol 71:398-405
 138. Lien RJ, Siopes TD 1993 The relationship of plasma thyroid hormone and prolactin concentrations to egg laying, incubation behavior, and molting by female turkeys exposed to a one-year natural daylength cycle. Gen Comp Endocrinol 90:205-213
 139. Takada M 1989 Effect of prolactin on the active Na transport across *Rana catesbeiana* skin during metamorphosis. Gen Comp Endocrinol 73:318-324
 140. Takada M 1986 The short-term effect of prolactin on the active Na transport system of the tadpole skin during metamorphosis. Comp Biochem Physiol 85:755-759
 141. Kern MD 1979 Seasonal changes in the reproductive system of the female white-crowned sparrow, *Zonotrichia leucophrys gambelii*, in captivity and in the field. 2. The incubation patch. Cell Tissue Res 202:379-398
 142. Sage M 1970 Control of prolactin release and its role in color change in the teleost *Gillichthys mirabilis*. J Exp Zool 173:121-127
 143. Duncan MJ, Goldman BD 1985 Physiological doses of prolactin stimulate pelage pigmentation in Djungarian hamster. Am J Physiol 248:664-667
 144. Girolomoni G, Phillips JT, Bergstresser PR 1993 Prolactin stimulates proliferation of cultured human keratinocytes. J Invest Dermatol 101:275-279
 145. Curlew JD, Loudon AS, Milne JA, McNeilly AS 1988 Effects of chronic long-term bromocriptine treatment on liveweight, voluntary food intake, coat growth and breeding season in non-pregnant red deer hinds. J Endocrinol 119:413-420
 146. Ibraheem M, Galbraith H, Scaife J, Ewen S 1994 Growth of secondary hair follicles of the Cashmere goat *in vitro* and their response to prolactin and melatonin. J Anat 185:135-142
 147. Dumont JN 1965 Prolactin-induced cytologic changes in the mucosa of the pigeon crop during crop-"milk" formation. Z Zellforsch Mikrosk Anat 68:755-782
 148. Anderson TR, Pitts DS, Nicoll CS 1984 Prolactin's mitogenic action on the pigeon crop-sac mucosal epithelium involves direct and indirect mechanisms. Gen Comp Endocrinol 54:236-246
 149. Hamosh M, Hamosh P 1977 The effect of prolactin on the lecithin content of fetal rabbit lung. J Clin Invest 59:1002-1005
 150. Hauth JC, Parker Jr CR, MacDonald PC, Porter JC, Johnston JM 1978 A role of fetal prolactin in lung maturation. Obstet Gynecol 51:81-88
 151. Quirk Jr JG, MacDonald PC, Johnston JM 1982 Role of fetal pituitary prolactin in fetal lung maturation. Semin Perinatol 6:238-245
 152. Johnson JW, Tyson JE, Mitzner W, Beck JC, Andreassen B, London WT, Villar J 1985 Amniotic fluid prolactin and fetal lung maturation. Am J Obstet Gynecol 153:372-380
 153. Richards JF 1975 Ornithine decarboxylase activity in tissues of prolactin-treated rats. Biochem Biophys Res Commun 63:292-299
 154. Vergani G, Mayerhofer A, Bartke A 1994 Acute effects of rat growth hormone (GH), human GH and prolactin on proliferating rat liver cells *in vitro*: a study of mitotic behaviour and ultrastructural alterations. Tissue Cell 26:457-465
 155. Buckley AR, Crowe PD, Russell DH 1988 Rapid activation of a protein kinase C in isolated rat liver nuclei by prolactin, a known hepatic mitogen. Proc Natl Acad Sci USA 85:8649-8653
 156. Crowe PD, Buckley AR, Zorn NE, Rui H 1991 Prolactin activates protein kinase C and stimulates growth-related gene expression in rat liver. Mol Cell Endocrinol 79:29-35
 157. Piccoletti R, Maroni P, Bendinelli P, Bernelli-Zazzera A 1994 Rapid stimulation of mitogen-activated protein kinase of rat liver by prolactin. Biochem J 303:429-433
 158. Santi P, Martelli AM, Gilmour RS, Falcieri E, Rana R, Cataldi A, Lattanzi G, Bareggi R, Cocco L 1992 Changes in phosphoinositide level in rat liver nuclei in response to prolactin, a known hepatic mitogen. Cell Signal 4:385-391
 159. Hill DJ, Francis MJ, Milner RD 1977 Action of rat prolactin on plasma somatomedin levels in the rat and on somatomedin release from perfused rat liver. J Endocrinol 75:137-143
 160. Ibrahim RH, Platt JE 1988 Enhancement of ornithine decarboxylase activity in *Ambystoma* liver slices by ovine prolactin: an evaluation of possible mediators. Gen Comp Endocrinol 72:90-96
 161. Hurley TW, Kuhn CM, Schanberg SM, Handwerker S 1980 Differential effects of placental lactogen, growth hormone and prolactin on rat liver ornithine decarboxylase activity in the perinatal period. Life Sci 27:2269-2275
 162. Berlanga JJ, Fresno Vara JA, Martin-Perez J, Garcia-Ruiz JP 1995 Prolactin receptor is associated with c-src kinase in rat liver. Mol Endocrinol 9:1461-1467
 163. Zhu XH, Zellweger R, Ayala A, Chaudry IH 1996 Prolactin inhibits the increased cytokine gene expression in Kupffer cells following haemorrhage. Cytokine 8:134-140
 164. Mick CC, Nicoll CS 1985 Prolactin directly stimulates the liver *in vivo* to secrete a factor (synlactin) which acts synergistically with the hormone. Endocrinology 116:2049-2053
 165. English DE, Russell SM, Katz LS, Nicoll CS 1990 Evidence for a role of the liver in the mammatrophic action of prolactin. Endocrinology 126:2252-2256
 166. Delidow BC, Hebert N, Steiny S, Nicoll CS 1986 Secretion of prolactin-synergizing activity (synlactin) by the liver of ectothermic vertebrates *in vitro*. J Exp Zool 238:147-153

167. Hoeffler JP, Frawley LS 1987 Liver tissue produces a potent lactogen that partially mimics the actions of prolactin. *Endocrinology* 120:1679-1681
168. Reddy PM, Reddy PR 1990 Effect of prolactin on DNA methylation in the liver and kidney of rat. *Mol Cell Biochem* 95:43-47
169. Bonga SE 1976 The effect of prolactin on kidney structure of the euryhaline teleost *Gasterosteus aculeatus* during adaptation to fresh water. *Cell Tissue Res* 166:319-338
170. Olivereau M, Lemoine AM 1969 Structural changes of the kidney in hypophysectomized cells treated with prolactin. *Z Zellforsch Mikrosk Anat* 95:361-376
171. Mainoya JR 1978 Possible influence of prolactin on intestinal hypertrophy in pregnant and lactating rats. *Experientia* 34:1230-1231
172. Muller Dowling RH 1981 Prolactin and the small intestine. Effect of hyperprolactinemia on mucosal structure in the rat. *Gut* 22:558-565
173. Goodlad RA, Writht NA 1990 Growth control factors in the gastrointestinal tract. *Baillieres Clin Gastroenterol* 4:97-118
174. Levi WM 1969 *The Pigeon*. Levi, Sumter, NC, pp 1-667
175. Bates RW, Miller RA, Garrison AM 1962 Evidence in the hypophysectomized pigeon of a synergism among prolactin, growth hormone, thyroxine and prednisone upon weight of the body, digestive tract, kidney and fat stores. *Endocrinology* 71:345-360
176. Sauro MD, Zorn NE 1991 Prolactin induces proliferation of vascular smooth muscle cells through a protein kinase C-dependent mechanism. *J Cell Physiol* 148:133-138
177. Sauro MD, Buckley AR, Russell DH, Fitzpatrick DF 1989 Prolactin stimulation of protein kinase C activity in rat aortic smooth muscle. *Life Sci* 44:1787-1792
178. Sauro MD, Bing B, Zorn NE 1992 Prolactin induces growth-related gene expression in rat aortic smooth muscle *in vivo*. *Eur J Pharmacol* 225:351-354
179. McAveney KM, Gimble JM, Yu-Lee L-Y 1996 Prolactin receptor expression during adipocyte differentiation of bone marrow stroma. *Endocrinology* 137:5723-5726
180. Brelje TC, Sorenson RL 1991 Role of prolactin vs. growth hormone on islet β -cell proliferation *in vitro*: implications for pregnancy. *Endocrinology* 128:45-57
181. Billestrup N, Nielsen JH 1991 The stimulatory effect of growth hormone, prolactin, and placental lactogen on β -cell proliferation is not mediated by insulin-like growth factor-I. *Endocrinology* 129:883-888
182. Nielsen JH, Moldrup A, Billestrup N, Petersen ED, Allevato G, Stahl M 1992 The role of growth hormone and prolactin in β cell growth and regeneration. *Adv Exp Med Biol* 321:9-17
183. Brelje TC, Parsons JA, Sorenson RL 1994 Regulation of islet β -cell proliferation by prolactin in rat islets. *Diabetes* 43:263-273
184. Markoff E, Beattie GM, Hayek A, Lewis UJ 1990 Effects of prolactin and glycosylated prolactin on (pro)insulin synthesis and insulin release from cultured rat pancreatic islets. *Pancreas* 5:99-103
185. Sorenson RL, Brelje TC, Hegre OD, Marshall S, Anaya P, Sheridan JD 1987 Prolactin (*in vitro*) decreases the glucose stimulation threshold, enhances insulin secretion, and increases dye coupling among islet β cells. *Endocrinology* 121:1447-1453
186. Michaels RL, Sorenson RL, Parsons JA, Sheridan JD 1987 Prolactin enhances cell-to-cell communication among β -cells in pancreatic islets. *Diabetes* 36:1098-1103
187. Krown KA, Wang YF, Ho TWC, Kelly PA, Walker AM 1992 Prolactin isoform 2 as an autocrine growth factor for GH₃ cells. *Endocrinology* 131:595-602
188. Yoshimura Y, Hosoi Y, Iritani A, Nakamura Y, Atlas SJ, Wallach EE 1989 Developmental potential of rabbit oocyte matured *in vitro*: the possible contribution of prolactin. *Biol Reprod* 41:26-33
189. Nag S, Sanyal S, Ghosh KK, Biswas NM 1981 Prolactin suppression and spermatogenic developments in maturing rats. A quantitative study. *Horm Res* 15:72-77
190. Buntin JD, Tesch D 1985 Effects of intracranial prolactin administration on maintenance of incubation readiness, ingestive behavior, and gonadal condition in ring doves. *Horm Behav* 19:188-203
191. Dombrowicz D, Sente B, Closset J, Hennen G 1992 Dose-dependent effects of human prolactin on the immature hypophysectomized rat testis. *Endocrinology* 130:695-700
192. Martinet L, Mondain-Monval M, Monnerie R 1992 Endogenous circannual rhythms and photorefractoriness of testis activity, moult and prolactin concentrations in mink (*Mustela vison*). *J Reprod Fertil* 95:325-338
193. Syms AJ, Harper ME, Griffiths K 1985 The effect of prolactin on human BPH epithelial cell proliferation. *Prostate* 6:145-153
194. Thomas JA, Manandhar MS, Keenan EJ, Edwards WD, Klase PA 1976 Effects of prolactin and dihydrotestosterone upon the rat prostate gland. *Urol Int* 31:265-271
195. Rui H, Purvis K 1987 Prolactin selectively stimulates ornithine decarboxylase in the lateral lobe of the rat prostate. *Mol Cell Endocrinol* 50:89-97
196. Negro-Vilar A, Saad WA, McCann SM 1977 Evidence for a role of prolactin in prostate and seminal vesicle growth in immature male rats. *Endocrinology* 100:729-737
197. Golander A, Weisman Y, Spierer Z, Binderman I, Kaye AM, Sömjén D 1988 Prolactin stimulates creatine kinase activity and DNA synthesis in explants of human amnion. *Acta Endocrinol (Copenh)* 119:223-227
198. DeVito WJ, Okulicz WC, Stone S, Avakian C 1992 Prolactin-stimulated mitogenesis of cultured astrocytes. *Endocrinology* 130:2549-2556
199. Shyr SW, Crowley WR, Grosvenor CE 1986 Effect of neonatal prolactin deficiency on prepubertal tuberoinfundibular and tuberohypophyseal dopaminergic neuronal activity. *Endocrinology* 119:1217-1221
200. Crim JW 1975 Prolactin-thyroxine antagonism and the metamorphosis of visual pigments in *Rana catesbeiana* tadpoles. *J Exp Zool* 192:355-362
201. Berczi I, Nagy E, de Toledo SM, Matusik RJ, Friesen HG 1991 Pituitary hormones regulate c-myc and DNA synthesis in lymphoid tissue. *J Immunol* 146:2201-2206
202. Hartmann DP, Holaday JW, Bernton EW 1989 Inhibition of lymphocyte proliferation by antibodies to prolactin. *FASEB J* 3:2194-2202
203. Mukherjee P, Mastro AM, Hymer WC 1990 Prolactin induction of interleukin-2 receptors on rat splenic lymphocytes. *Endocrinology* 126:88-94
204. Bhatavdekar J, Patel DD, Ghosh N, Vora H, Shah N, Karelia N, Balar D, Chikhlikar P, Dave R 1994 Interrelationship of prolactin and its receptor in carcinoma of colon and rectum: a preliminary report. *J Surg Oncol* 55:246-249
205. Bhatavdekar JM, Patel DD, Shah NG, Karelia NH, Vora HH, Ghosh N, Suthar TP, Balar DB 1995 Prognostic value of insulin-like growth factor-1 receptors in patients with colon/rectal cancer: correlation with plasma prolactin. *Eur J Surg Oncol* 21:23-26
206. Kiss R, de Launoit Y, L'Hermite-Balériaux M, L'Hermite M, Paridaens RJ, Danguy AJ, Pasteels JL 1987 Effect of prolactin and estradiol on cell proliferation in the uterus and the MXT mouse mammary neoplasm. *J Natl Cancer Inst* 78:993-998
207. Biswas R, Vonderhaar BK 1987 Role of serum in the prolactin responsiveness of MCF-7 human breast cancer cells in long-term tissue culture. *Cancer Res* 47:3509-3514
208. Das R, Vonderhaar BK 1996 Involvement of SHC, GRB2, SOS and RAS in prolactin signal transduction in mammary epithelial cells. *Oncogene* 13:1139-1145
209. Walker AM, Montgomery DW, Saraiya S, Ho TW, Garewal HS, Wilson J 1995 Prolactin-immunoglobulin G complexes from human serum act as costimulatory ligands causing proliferation of malignant B lymphocytes. *Proc Natl Acad Sci USA* 92:3278-3282
210. Barker CS, Bear SE, Keler T, Copeland NG, Gilbert DJ, Jenkins NA, Yeung RS, Tschlis PN 1992 Activation of the prolactin receptor gene by promoter insertion in a moloney murine leukemia virus-induced rat thymoma. *J Virol* 66:6763-6768
211. Gout PW, Beer CT, Noble RL 1980 Prolactin-stimulated growth of cell cultures established from malignant Nb rat lymphomas. *Cancer Res* 40:2433-2436
212. Nishiguchi Y, Hibasami H, Komada Y, Sakurai M, Nakashima K 1993 Human promyelocytic cell line HL60 has the specific binding sites for prolactin and its ornithine decarboxylase, DNA synthesis and cellular proliferation are induced by prolactin. *Leuk Res* 17:633-637
213. Nowak RA, Rein MS, Heffner LJ, Friedman AJ, Tashjian AH 1993

- Production of prolactin by smooth muscle cells cultured from human uterine fibroid tumors. *J Clin Endocrinol Metab* 76:1308-1313
214. Ormandy CJ, Camus A, Barra J, Damotte D, Lucas BK, Buteau H, Edery M, Brousse N, Babinet C, Binart N, Kelly PA 1997 Null mutation of the prolactin receptor gene produces multiple reproductive defects in the mouse. *Genes Dev* 11:167-178
 215. Brown-Borg HM, Borg KE, Meliska CJ, Bartke A 1996 Dwarf mice and the ageing process. *Nature* 384:33
 216. Kumaran B, Gunasekar PG, Aruldas MM, Govindarajulu P 1989 Influence of prolactin on neural and glial cellular adenosine triphosphatases in immature male bonnet monkeys (*Macaca radiata*, Geoffroy). *Indian J Exp Biol* 27:17-19
 217. Machida T, Taga M, Minaguchi H 1990 Effect of prolactin (PRL) on lipoprotein lipase (LPL) activity in the rat fetal liver. *Asia Oceania J Obstet Gynaecol* 16:261-265
 218. Liu Y, Hyde JF, Vore M 1992 Prolactin regulates maternal bile secretory function post partum. *J Pharmacol Exp Ther* 261:560-566
 219. Ganguly TC, Liu Y, Hyde JF, Hagenbuch B, Meier PJ, Vore M 1994 Prolactin increases hepatic Na⁺/taurocholate co-transport activity and messenger RNA post partum. *Biochem J* 303:33-36
 220. Liu Y, Ganguly T, Hyde JF, Vore M 1995 Prolactin increases mRNA encoding Na⁺-TC co-transport polypeptide and hepatic Na⁺-TC co-transport. *Am J Physiol* 268:11-17
 221. Garrison MM, Scow RO 1975 Effect of prolactin on lipoprotein lipase in crop sac and adipose tissue of pigeons. *Am J Physiol* 228:1542-1544
 222. Kumaran B, Gunasekar PG, Aruldas MM, Govindarajulu P 1988 Role of prolactin on neural and glial cellular enzymes involved in carbohydrate metabolism. I. Studies on immature male bonnet monkeys. *Brain Res* 450:325-333
 223. Villalba M, Zabala MT, Martinez-Serrano A, de la Colina R, Satrustegui J, Garcia-Ruiz JP 1991 Prolactin increases cytosolic free calcium concentration in hepatocytes of lactating rats. *Endocrinology* 129:2857-2861
 224. Curry DL, Bennett LL, Li CH 1975 Dynamics of insulin release by perfused hamster (*Mesocricetus auratus*) pancreases: effects of hypophysectomy, bovine and human growth hormone, and prolactin. *J Endocrinol* 65:245-251
 225. Nielsen JH 1982 Effects of growth hormone, prolactin, and placental lactogen on insulin content and release, and deoxyribonucleic acid synthesis in cultured pancreatic islets. *Endocrinology* 110:600-606
 226. Sorenson RL, Johnson MG, Parsons JA, Sheridan JD 1987 Decreased glucose stimulation threshold, enhanced insulin secretion, and increased β -cell coupling in islets of prolactin-treated rats. *Pancreas* 2:283-288
 227. Weinhaus AJ, Stout LE, Sorenson RL 1996 Glucokinase, hexokinase, glucose transporter 2, and glucose metabolism in islets during pregnancy and prolactin-treated islets *in vitro*: mechanisms for long term up-regulation of islets. *Endocrinology* 137:1640-1649
 228. Glasow A, Breidert M, Haidan A, Anderegg U, Kelly PA, Bornstein S 1996 Functional aspects of the effect of prolactin (PRL) on adrenal steroidogenesis and distribution of the PRL receptor in the human adrenal gland. *J Clin Endocrinol Metab* 81:3103-3111
 229. Higuchi K, Nawata H, Maki T, Higashizima M, Kato K, Ibayashi H 1984 Prolactin has a direct effect on adrenal androgen secretion. *J Clin Endocrinol Metab* 59:714-718
 230. Albertson BD, Sienkiewicz ML, Kimball D, Munabi AK, Cassorla F, Loriaux DL 1987 New evidence for a direct effect of prolactin on rat adrenal steroidogenesis. *Endocr Res* 13:317-333
 231. Couet J, Martel C, Labrie Y, Luo S, Simard J, Labrie F 1994 Opposite effects of prolactin and corticosterone on the expression and activity of 3 β -hydroxysteroid dehydrogenase/ Δ^5 - Δ^4 isomerase in rat skin. *J Invest Dermatol* 103:60-64
 232. Dave JR, Brown NV, Knazek RA 1982 Prolactin modifies the prostaglandin synthesis, prolactin binding and fluidity of mouse liver membranes. *Biochem Biophys Res Commun* 108:193-199
 233. Murphy IJ, Tachibana K, Friesen HG 1988 Stimulation of hepatic insulin-like growth factor-I gene expression by ovine prolactin: evidence for intrinsic somatogenic activity in the rat. *Endocrinology* 122:2027-2033
 234. Fernandez-Ruiz JJ, Ceberia M, Agrasal C, Tresguerres JAF, Esquifino AI, Ramos JA 1987 Effect of elevated prolactin levels on the synthesis and release of catecholamines from the adrenal medulla in female rats. *Neuroendocrinology* 45:208-211
 235. Fernandez-Ruiz JJ, Martinez-Arrieta R, Hernandez ML, Ramos JA 1988 Possible direct effect of prolactin on catecholamine synthesis and release in rat adrenal medulla: *in vitro* studies. *J Endocrinol Invest* 11:603-608
 236. Blüm V, Fiedler K 1965 Hormonal control of reproductive behavior in some cichlid fish. *Gen Comp Endocrinol* 5:186-196
 237. Machemer L, Fiedler K 1965 Zur hormonalen steuerung des schaumnestbaues beim paradiesfisch, *Macropodus opercularis* L. (Anabantidae, Teleostei). *Naturwissenschaften* 52:648-649
 238. Lam TJ, Hoar WS 1967 Seasonal effects of prolactin on fresh-water osmoregulation of the marine form (trachurus) of the stickleback *Gasterosteus aculeatus*. *Can J Zool* 45:509-516
 239. Meier AH, Farmer DS, King JR 1965 A possible endocrine basis for migratory behavior in the white-crowned sparrow, *Zonotrichia leucophrys gambelii*. *Anim Behav* 13:453-465
 240. Leboucher G, Richard-Yris MA, Williams J, Chadwick A 1990 Incubation and maternal behaviour in domestic hens: influence of the presence of chicks on circulating luteinising hormone, prolactin and oestradiol and on behaviour. *Br Poult Sci* 31:851-862
 241. Horseman ND, Buntin JD 1995 Regulation of pigeon cropmilk secretion and parental behaviors by prolactin. *Annu Rev Nutr* 15:213-238
 242. Janik DS, Buntin JD 1985 Behavioural and physiological effects of prolactin in incubating ring doves. *J Endocrinol* 105:201-209
 243. Hector JA, Goldsmith AR 1985 The role of prolactin during incubation: comparative studies of three Diomedea albatrosses. *Gen Comp Endocrinol* 60:236-243
 244. March JB, Sharp PJ, Wilson PW, Sang HM 1994 Effect of active immunization against recombinant-derived chicken prolactin fusion protein on the onset of broodiness and photoinduced egg laying in bantam hens. *J Reprod Fertil* 101:227-233
 245. Bridges RS, Dibiase R, Loundes DD, Doherty PC 1985 Prolactin stimulation of maternal behavior in female rats. *Science* 227:782-784
 246. Bridges RS, Numan M, Ronsheim PM, Mann PE, Lupini CE 1990 Central prolactin infusions stimulate maternal behavior in steroid-treated, nulliparous female rats. *Proc Natl Acad Sci USA* 87:8003-8007
 247. Bridges RS, Robertson MC, Shiu RP, Sturgis JD, Henriquez BM, Mann PE 1997 Central lactogenic regulation of maternal behavior in rats: steroid dependence, hormone specificity, and behavioral potencies of rat prolactin and rat placental lactogen I. *Endocrinology* 138:756-763
 248. Drago F, Canonico PL, Bitetti R, Scapagnini U 1980 Systemic and intraventricular prolactin induces excessive grooming. *Eur J Pharmacol* 65:457-458
 249. Drago F, Bohus B, Gispen WH, Scapagnini U, De Wied D 1983 Prolactin-enhanced grooming behavior: interaction with ACTH. *Brain Res* 263:277-282
 250. Drago F, D'Agata V, Iacona T, Spadaro F, Grassi M, Valerio C, Raffaele R, Astuto C, Lauria N, Vitetta M 1989 Prolactin as a protective factor in stress-induced biological changes. *J Clin Lab Anal* 3:340-344
 251. Das K 1991 Effects of testosterone propionate, prolactin and photoperiod on feeding behaviours of Indian male weaver birds. *Indian J Exp Biol* 29:1104-1108
 252. Gerardo-Gettens T, Moore BJ, Stern JS, Horwitz BA 1989 Prolactin stimulates food intake in a dose-dependent manner. *Am J Physiol* 256:276-280
 253. Noel MB, Woodside B 1993 Effects of systemic and central prolactin injections on food intake, weight gain, and estrous cyclicity in female rats. *Physiol Behav* 54:151-154
 254. Buntin JD, Becker GM, Ruzycski E 1991 Facilitation of parental behavior in ring doves by systemic or intracranial injections of prolactin. *Horm Behav* 25:424-444
 255. Alvarez EO, Banzan AM 1994 Behavioural actions of prolactin locally applied into the hippocampus of adult female rats. *J Neural Transm* 95:17-28
 256. Nicoletti F, Drago F, Speciale C, Kovacs LG, Scapagnini U 1983 Role of prolactin-opiate interactions in the central regulation of pain threshold. *Cephalalgia* 3:31-34

257. Ramaswamy S, Viswanathan S, Bapna JS 1987 Effect of prolactin on tolerance and dependence to acute administration of morphine. *Neuropharmacology* 26:111-113
258. Ramaswamy S, Pillai NP, Bapna JS 1983 Analgesic effect of prolactin: possible mechanism of action. *Eur J Pharmacol* 96:171-173
259. Ramaswamy S, Suthakaran C, Bapna JS 1989 Role of GABAergic system in prolactin analgesia. *Clin Exp Pharmacol Physiol* 16:893-896
260. Ramaswamy S, Regunathan S, Bapna JS 1988 Anticholinesterase activity of prolactin: correlation with analgesia. *J Pharm Pharmacol* 40:306-307
261. Ramaswamy S, Rajasekaran M, Bapna JS 1986 Role of calcium in prolactin analgesia. *Arch Int Pharmacodyn Ther* 283:56-60
262. Shewade DG, Ramaswamy S 1995 Prolactin induced analgesia is dependent on ATP sensitive potassium channels. *Clin Exp Pharmacol Physiol* 22:635-636
263. Sobrinho LG 1993 The psychogenic effects of prolactin. *Acta Endocrinol (Copenh)* 129:38-40
264. Howles CM, Webster GM, Haynes NB 1980 The effect of rearing under a long or short photoperiod on testis growth, plasma testosterone and prolactin concentrations, and the development of sexual behaviour in rams. *J Reprod Fertil* 60:437-447
265. Obal Jr F, Opp M, Cady AB, Johannsen L, Krueger JM 1989 Prolactin, vasoactive intestinal peptide, and peptide histidine methionine elicit selective increases in REM sleep in rabbits. *Brain Res* 490:292-300
266. Obal Jr F, Payne L, Kacsoh B, Opp M, Kapas L, Grosvenor CE, Krueger JM 1994 Involvement of prolactin in the REM sleep-promoting activity of systemic vasoactive intestinal peptide (VIP). *Brain Res* 645:143-149
267. Roky R, Obal Jr F, Valatx JL, Bredow S, Fang J, Pagano LP, Krueger JM 1995 Prolactin and rapid eye movement sleep regulation. *Sleep* 18:536-542
268. Roky R, Valatx JL, Jouvett M 1993 Effect of prolactin on the sleep-wake cycle in the rat. *Neurosci Lett* 156:117-120
269. Roky R, Valatx JL, Paut-Pagano L, Jouvett M 1994 Hypothalamic injection of prolactin or its antibody alters the rat sleep-wake cycle. *Physiol Behav* 55:1015-1019
270. Harlan RE, Shivers BD, Pfaff DW 1983 Midbrain microinfusions of prolactin increase the estrogen-dependent behavior, lordosis. *Science* 219:1451-1453
271. Lookingland KJ, Moore KE 1984 Effects of estradiol and prolactin on incertohypothalamic dopaminergic neurons in the male rat. *Brain Res* 323:83-91
272. Moore KE, Demarest KT 1982 Effects of prolactin on central dopaminergic neurons. *Prog Neuropsychopharmacol Biol Psychiatry* 6:645-650
273. Silverman WF, Walsh RJ 1987 Prolactin causes increased turnover of dopamine in 10-day-old rat median eminence. *Anat Rec* 217:53-55
274. Chen JC, Ramirez VD 1989 Effects of prolactin on tyrosine hydroxylase activity of central dopaminergic neurons of male rats. *Eur J Pharmacol* 166:473-479
275. Shah GV, Shyr SW, Grosvenor CE, Crowley WR 1988 Hyperprolactinemia after neonatal prolactin (PRL) deficiency in rats: evidence for altered anterior pituitary regulation of PRL secretion. *Endocrinology* 122:1883-1889
276. Voogt JL, de Greef WJ, Visser TJ, de Koning J, Vreeburg JTM, Weber RF 1987 *In vivo* release of dopamine, luteinizing hormone-releasing hormone and thyrotropin-releasing hormone in male rats bearing a prolactin-secreting tumor. *Neuroendocrinology* 46:110-116
277. Sarkar DK, Yen SS 1985 Hyperprolactinemia decreases the luteinizing hormone-releasing hormone concentration in pituitary portal plasma: a possible role for β -endorphin as a mediator. *Endocrinology* 116:2080-2084
278. Cohen-Becker IR, Selmanoff M, Wise PM 1986 Hyperprolactinemia alters the frequency and amplitude of pulsatile luteinizing hormone secretion in the ovariectomized rat. *Neuroendocrinology* 42:328-333
279. Chan A, Dudley CA, Moss RL 1983 Action of prolactin, dopamine and LHRH on ventromedial hypothalamic neurons as a function of ovarian hormones. *Neuroendocrinology* 36:397-403
280. DeVito WJ, Stone S, Avakian C 1991 Prolactin stimulation of protein kinase C activity in the rat hypothalamus. *Biochem Biophys Res Commun* 176:660-667
281. Sharif NA 1988 Chronic prolactin, gonadal and thyroid hormone treatments *in vivo* alter levels of TRH and muscarinic receptors in male and female rat tissues. *Brain Res* 449:364-368
282. O'Steen WK, Kraeer SL 1977 Effects of hypophysectomy, pituitary gland homogenates and transplants, and prolactin on photoreceptor destruction. Association for Research in Vision and Ophthalmology. *Invest Ophthalmol Vis Sci*, 16:940-946
283. Grosvenor CE, Whitworth NS 1983 Accumulation of prolactin in maternal milk and its transfer to circulation of neonate rat: a review. *Endocrinol Exp* 17:271-282
284. Neville MC, Daniel CW (eds) 1987 *The Mammary Gland: Development, Regulation and Function*. Plenum Press, New York
285. Ono M, Oka T 1980 The differential actions of cortisol on the accumulation of α lactalbumin and casein in midpregnant mouse mammary gland in culture. *Cell* 19:473-480
286. Guyette WA, Matusik RJ, Rosen JM 1979 Prolactin-mediated transcriptional and post-transcriptional control of casein gene expression. *Cell* 17:1013-1023
287. Hennighausen L, Westphal C, Sankaran L, Pittius CW 1991 Regulation of expression of genes for milk proteins. *Biotechnology* 16:65-74
288. Rosen JM, Rodgers JR, Couch CH, Bisbee CA, David-Inouye Y, Campbell SM, Yu-Lee L-Y 1986 Multihormonal regulation of milk protein gene expression. *Ann NY Acad Sci* 478:63-76
289. Collet C, Joseph R, Nicholas K 1991 A marsupial β -lactoglobulin gene-characterization and prolactin-dependent expression. *J Mol Endocrinol* 6:9-16
290. Oppat CA, Rillema JA 1988 Characteristics of the early effect of prolactin on lactose biosynthesis in mouse mammary gland explants. *Proc Soc Exp Biol Med* 188:342-345
291. Jagoda CA, Rillema JA 1991 Temporal effect of prolactin on the activities of lactose synthetase, α -lactalbumin, and galactosyl transferase in mouse mammary gland explants. *Proc Soc Exp Biol Med* 197:431-434
292. Waters SB, Rillema JA 1988 Effect of prolactin on enzymes of lipid biosynthesis in mammary gland explants. *Am J Physiol* 255:E567-E571
293. Barber MC, Travers MT, Finley E, Flint DJ, Vernon RG 1992 Growth-hormone prolactin interactions in the regulation of mammary and adipose-tissue acetyl-coA carboxylase activity and gene expression in lactating rats. *Biochem J* 285:469-475
294. Barber MC, Finley E, Vernon RG 1991 Mechanisms whereby prolactin modulates lipogenesis in sheep mammary gland. *Horm Metab Res* 23:143-145
295. Barber MC, Clegg RA, Finley E, Vernon RG, Flint DJ 1992 The role of growth hormone, prolactin and insulin-like growth factors in the regulation of rat mammary gland and adipose tissue metabolism during lactation. *J Endocrinol* 135:195-202
296. Fielder PJ, Thordarson G, English A, Rosenfeld RG, Talamantes F 1992 Expression of a lactogen-dependent insulin-like growth factor-binding protein in cultured mouse mammary epithelial cells. *Endocrinology* 131:261-267
297. Fenton SE, Sheffield LG 1991 Lactogenic hormones increase epidermal growth factor messenger RNA content of mouse mammary glands. *Biochem Biophys Res Commun* 181:1063-1069
298. Ball RK, Friis RR, Schoenenberger CA, Groner B 1988 Prolactin regulation of β -casein gene expression and of a cytosolic 120-kd protein in a cloned mouse mammary epithelial cell line. *EMBO J* 7:2089-2095
299. Parry G, Stubbs J, Bissell MJ, Schmidhauser C, Spicer AP, Gendler SJ 1992 Studies of Muc-1 mucin expression and polarity in the mouse mammary gland demonstrate developmental regulation of Muc-1 glycosylation and establish the hormonal basis for mRNA expression. *J Cell Sci* 101:191-199
300. Shiu RP, Iwaszow BM 1985 Prolactin-inducible proteins in human breast cancer cells. *J Biol Chem* 260:11307-11313
301. Thiede MA, Rodan GA 1988 Expression of a calcium-mobilizing parathyroid hormone-like peptide in lactating mammary tissue. *Science* 242:278-280
302. Horseman ND 1989 A prolactin-inducible gene product which is

- a member of the calpactin/lipocortin family. *Mol Endocrinol* 3:773-779
303. Pukac LA, Horseman ND 1984 Regulation of pigeon crop gene expression by prolactin. *Endocrinology* 114:1718-1724
 304. Horseman ND, Will CL 1984 Characterization of lipid storage associated proteins induced in crop tissue by prolactin. *J Comp Physiol B* 154:237-242
 305. Horseman ND, Nollin LJ 1985 The mitogenic, but not differentiative, response of crop tissue to prolactin is circadian phase dependent. *Endocrinology* 116:2085-2089
 306. Pukac LA, Horseman ND 1987 Regulation of cloned prolactin-inducible genes in pigeon crop. *Mol Endocrinol* 1:188-194
 307. Armstrong DT, Knudsen KA, Miller LS 1970 Effects of prolactin upon cholesterol metabolism and progesterone biosynthesis in corpora lutea of rats hypophysectomized during pseudopregnancy. *Endocrinology* 86:634-641
 308. Ota H, Wakizaka A, Fukushima M, Maki M 1985 Dual regulation of rat ovarian LH-receptor by the administration of prolactin or sulpiride. *IRCS J Med Sci* 63:257-264
 309. Berhman HR, Orczyk GP, MacDonald GJ, Greep RO 1970 Prolactin induction of enzymes controlling luteal cholesterol ester turnover. *Endocrinology* 87:1251-1256
 310. Dunaif AE, Zimmerman EA, Friesen HG, Frantz AG 1982 Intracellular localization of prolactin receptor and prolactin in the rat ovary by immunocytochemistry. *Endocrinology* 110:1465-1471
 311. Larsen JL, Bhanu A, Odell WD 1990 Prolactin inhibition of pregnant mare's serum stimulated follicle development in the rat ovary. *Endocr Res* 16:449-459
 312. Martel C, Labrie C, Dupont E, Couet J, Trudel C, Rheume E, Simard J, Luu-The V, Pelletier G, Labrie F 1990 Regulation of 3 β -hydroxysteroid dehydrogenase/ Δ_5 - Δ_4 isomerase expression and activity in the hypophysectomized rat ovary: interactions between the stimulatory effect of human chorionic gonadotropin and the luteolytic effect of prolactin. *Endocrinology* 127:2726-2737
 313. Tsai-Morris CH, Ghosh M, Hirshfield AN, Wise PM, Brodie AM 1983 Inhibition of ovarian aromatase by prolactin *in vivo*. *Biol Reprod* 29:342-346
 314. Martel C, Labrie C, Couet J, Dupont E, Trudel C, Luu-The V, Takahashi M, Pelletier G, Labrie F 1990 Effects of human chorionic gonadotropin (hCG) and prolactin (PRL) on 3 β -hydroxy-5-ene-steroid dehydrogenase/ Δ_5 - Δ_4 isomerase (3 β -HSD) expression and activity in the rat ovary. *Mol Cell Endocrinol* 18208:19975-16715
 315. Yoshimura Y, Tada S, Oda T, Nakamura Y, Maruyama K, Ichikawa F, Ebihara T, Hirota Y, Sawada T, Kawakami S 1989 Direct inhibitory ovarian effects of prolactin in the process of ovulation. *Nippon Sanka Fujinka Gakkaishi* 41:83-89
 316. Zadworny D, Shimada K, Ishida H, Sato K 1989 Gonadotropin-stimulated estradiol production in small ovarian follicles of the hen is suppressed by physiological concentrations of prolactin *in vitro*. *Gen Comp Endocrinol* 74:468-473
 317. Yoshimura Y, Nakamura Y, Oda T, Ando M, Ubukata Y, Koyama N, Karube M, Yamada H 1992 Effects of prolactin on ovarian plasmin generation in the process of ovulation. *Biol Reprod* 46:322-327
 318. Yoshimura Y, Maruyama K, Shiraki M, Kawakami S, Fukushima M, Nakamura Y 1990 Prolactin inhibits plasminogen activator activity in the preovulatory follicles. *Endocrinology* 126:631-636
 319. Wang C, Hsueh AJ, Erickson GF 1980 Prolactin inhibition of estrogen production by cultured rat granulosa cells. *Mol Cell Endocrinol* 20:135-144
 320. Wang C, Chan V 1982 Divergent effects of prolactin on estrogen and progesterone production by granulosa cells of rat Graafian follicles. *Endocrinology* 110:1085-1093
 321. Fortune JE, Wissler RN, Vincent SE 1986 Prolactin modulates steroidogenesis by rat granulosa cells: II. Effects on estradiol. *Biol Reprod* 35:92-99
 322. Krasnow JS, Hickey GJ, Richards JS 1990 Regulation of aromatase mRNA and estradiol biosynthesis in rat ovarian granulosa and luteal cells by prolactin. *Mol Endocrinol* 4:13-22
 323. Rajkumar K, Klingshorn P, Chedrese PJ, Murphy BD 1988 Prolactin, LH and estradiol-17 β in utilization of lipoprotein substrate by porcine granulosa cells *in vitro*. *Can J Physiol Pharmacol* 66:561-566
 324. Chedrese PJ, Rajkumar K, Ly H, Murphy BD 1988 Dose response of luteinized porcine granulosa cells *in vitro* to prolactin: dependency on pre-exposure to human chorionic gonadotropin. *Can J Physiol Pharmacol* 66:1337-1340
 325. Jones PB, Valk CA, Hsueh AJ 1983 Regulation of progesterone biosynthetic enzymes in cultured rat granulosa cells: effects of prolactin, β_2 -adrenergic agonist, human chorionic gonadotropin and gonadotropin releasing hormone. *Biol Reprod* 29:572-585
 326. Casper RF, Erickson GF 1981 *In vitro* heteroregulation of LH receptors by prolactin and FSH in rat granulosa cells. *Mol Cell Endocrinol* 23:161-171
 327. Polan ML, Laufer N, Dlugi AM, Tarlatzis BC, Haseltine FP, DeCherney AH, Behrman HR 1984 Human chorionic gonadotropin and prolactin modulation of early luteal function and luteinizing hormone receptor-binding activity in cultured human granulosa-luteal cells. *J Clin Endocrinol Metab* 59:773-779
 328. Soto EA, Tureck RW, Strauss III JF 1985 Effects of prolactin on progesterone secretion by human granulosa cells in culture. *Biol Reprod* 32:541-545
 329. Cutie E, Andino NA 1988 Prolactin inhibits the steroidogenesis in midfollicular phase human granulosa cells cultured in a chemically defined medium. *Fertil Steril* 49:632-637
 330. Lee MS, Ben-Rafael Z, Meloni F, Mastroianni Jr L, Flickinger GL 1986 Effects of prolactin on steroidogenesis by human luteinized granulosa cells. *Fertil Steril* 46:32-36
 331. Adashi EY, Resnick CE 1987 Prolactin as an inhibitor of granulosa cell luteinization: implications for hyperprolactinemia-associated luteal phase dysfunction. *Fertil Steril* 48:131-139
 332. Gitay-Goren H, Lindenbaum ES, Kraiem Z 1989 Prolactin inhibits hCG-stimulated steroidogenesis and cAMP accumulation, possibly by increasing phosphodiesterase activity, in rat granulosa cell cultures. *Mol Cell Endocrinol* 61:69-76
 333. Gitay-Goren H, Kraiem Z, Lindenbaum ES 1988 Effect of prolactin on the morphology of cultured rat granulosa cells. *Cytobios* 56:89-99
 334. Yamanouchi K, Matsuyama S, Nishihara M, Shiota K, Tachi C, Takahashi M 1992 Splenic macrophages enhance prolactin-induced progesterone secretion from mature rat granulosa cells *in vitro*. *Biol Reprod* 46:1109-1113
 335. Fanjul LF, Marrero I, Gonzales J, Quintana J, Santana P, Estévez F, Mato JM, Ruiz de Galarreta CM 1993 Does oligosaccharide-phosphatidylinositol (glycosyl-phosphatidylinositol) hydrolysis mediate prolactin signal transduction in granulosa cells? *Eur J Biochem* 216:747-755
 336. Dajee M, Kazansky AV, Raught B, Hocker G, Fey GH, Richards JS 1996 Prolactin induction of the α_2 -macroglobulin gene in rat ovarian granulosa cells: Stat5 activation and binding to the interleukin-6 response element. *Mol Endocrinol* 10:171-184
 337. Yoshimura Y, Nakamura Y, Yamada H, Ando M, Ubukata Y, Oda T, Suzuki M 1991 Possible contribution of prolactin in the process of ovulation and oocyte maturation. *Horm Res* 35:22-32
 338. Cecim M, Kerr J, Bartke A 1995 Infertility in transgenic mice overexpressing the bovine growth hormone gene: luteal failure secondary to prolactin deficiency. *Biol Reprod* 52:1162-1166
 339. Loudon AS, Brinklow BR, Gulland FD, Boyle J, Flint AP 1990 Roles of prolactin and the uterus in the control of luteal regression in the Bennett's wallaby (*Macropus rufogriseus rufogriseus*). *Reprod Fertil Dev* 2:71-78
 340. Wüttke W, Meites J 1971 Luteolytic role of prolactin during the estrous cycle of the rat. *Proc Soc Exp Biol Med* 137:988-991
 341. Matsuyama S, Shiota K, Takahashi M 1990 Possible role of transforming growth factor- β as a mediator of luteotropic action of prolactin in rat luteal cell cultures. *Endocrinology* 127:1561-1567
 342. Bernard RT, Bojarski C 1994 Effects of prolactin and hCG treatment on luteal activity and the conceptus during delayed implantation in Schreibers' long-fingered bat (*Miniopterus schreibersii*). *J Reprod Fertil* 100:359-365
 343. Albarracín CT, Gibori G 1991 Prolactin action on luteal protein expression in the corpus luteum. *Endocrinology* 129:1821-1830
 344. Daniel Jr JC, Jetton AE, Chilton BS 1984 Prolactin as a factor in the uterine response to progesterone in rabbits. *J Reprod Fertil* 72:443-452
 345. Chilton BS, Daniel Jr JC 1987 Differences in the rabbit uterine

- response to progesterone as influenced by growth hormone or prolactin. *J Reprod Fertil* 79:581-587
346. Prigent-Tessier A, Pageaux JF, Fayard JM, Lagarde M, Laugier C, Cohen H 1996 Prolactin up-regulates prostaglandin E₂ production through increased expression of pancreatic-type phospholipase A₂ (type I) and prostaglandin G/H synthase 2 in uterine cells. *Mol Cell Endocrinol* 122:101-108
 347. Kennedy TG, Armstrong DT 1972 Extra-ovarian effect of prolactin on vaginal mucification in the rat. *Endocrinology* 90:815-822
 348. Armstrong DT, King ER 1971 Uterine progesterone metabolism and progestational response: effects of estrogens and prolactin. *Endocrinology* 89:191-197
 349. Leung BS, Sasaki GH 1973 Prolactin and progesterone effect on specific estradiol binding in uterine and mammary tissues *in vitro*. *Biochem Biophys Res Commun* 55:1180-1187
 350. Saiduddin S, Zassenhaus HP 1977 Effect of prolactin on specific oestradiol receptors in the rat uterus. *J Endocrinol* 72:101-102
 351. Mati JK, Mugambi M, Muriuki PB, Thairu K 1974 Effect of prolactin on isolated rabbit myometrium. *J Endocrinol* 60:379-380
 352. Chilton BS, Mani SK, Bullock DW 1988 Servomechanism of prolactin and progesterone in regulating uterine gene expression. *Mol Endocrinol* 2:1169-1175
 353. Daniel Jr JC, Juneja SC, Taylor SP, Lonergan PB, Sullivan PK, Chilton BS 1988 Variability in the response of the rabbit uterus to progesterone as influenced by prolactin. *J Reprod Fertil* 84:13-21
 354. Randall GW, Daniel Jr JC, Chilton BS 1991 Prolactin enhances uteroglobin gene expression by uteri of immature rabbits. *J Reprod Fertil* 91:249-257
 355. Vijayan E, Jayashree J 1993 Prolactin suppression during pre- and post-implantation periods on rat uterine glucosamine synthase activity. *Indian J Exp Biol* 31:386-388
 356. Purvis K, Clausen OP, Olsen A, Haug E, Hansson V 1979 Prolactin and Leydig cell responsiveness to LH/hCG in the rat. *Arch Androl* 3:219-230
 357. Papadopoulos V, Drosowsky MA, Carreau S 1986 *In vitro* effects of prolactin and dexamethasone on rat Leydig cell aromatase activity. *Andrologia* 18:79-83
 358. Mathur U, Bartke A, Weisz J 1975 Effects of prolactin and LH on the activity of Δ^5 - 3β hydroxy-steroid dehydrogenase, dihydro-otrotic dehydrogenase, β -hydroxybutyrate dehydrogenase and glucose-6-phosphate dehydrogenase in the testis of the dwarf mice. *Indian J Physiol Pharmacol* 19:58-64
 359. Rubin RT, Poland RE, Tower BB 1976 Prolactin-related testosterone secretion in normal adult men. *J Clin Endocrinol Metab* 42:112-116
 360. Gunasekar PG, Kumaran B, Govindarajulu P 1988 Prolactin and Leydig cell steroidogenic enzymes in the bonnet monkey (*Macaca radiata*). *Int J Androl* 11:53-59
 361. Guillaumot P, Tabone E, Benahmed M 1996 Sertoli cells as potential targets of prolactin action in the testis. *Mol Cell Endocrinol* 122:199-206
 362. Gunasekar PG, Kumaran B, Govindarajulu P 1991 Role of prolactin on Leydig, Sertoli and germ cellular neutral lipids in bonnet monkeys, *Macaca radiata*. *Endocrinol Jpn* 38:1-8
 363. Reyes A, Parra A, Chavarria ME, Goicoechea B, Rosado A 1979 Effect of prolactin on the calcium binding and/or transport of ejaculated and epididymal human spermatozoa. *Fertil Steril* 31:669-672
 364. Pedron N, Giner J 1978 Effect of prolactin on the glycolytic metabolism of spermatozoa from infertile subjects. *Fertil Steril* 29:428-430
 365. Sheth AR, Gunjekar AN, Shah GV 1979 Effect of LH, prolactin and spermine on ATPase activity of human spermatozoa. *Andrologia* 11:11-14
 366. Shah GV, Desai RB, Sheth AR 1976 Effect of prolactin on metabolism of human spermatozoa. *Fertil Steril* 27:1292-1294
 367. Velazquez-Ramirez A, Vilar-Rojas C, Hicks JJ 1980 Similar effects of prolactin and dbcAMP upon human spermatozoa metabolism. *Int J Androl* 3:23-31
 368. Shah GV, Gunjekar AN, Sheth AR, Raut SJ 1980 Effect of prolactin and spermine on the zinc content of human spermatozoa. *Andrologia* 12:207-210
 369. Fukuda A, Mori C, Hashimoto H, Noda Y, Mori T, Hoshino K 1989 Effects of prolactin during preincubation of mouse spermatozoa on fertilizing capacity *in vitro*. *J In Vitro Fert Embryo Transfer* 6:92-97
 370. Reddy YD, Reddy KV, Govindappa S 1985 Effect of prolactin and bromocriptine administration on epididymal function: a biochemical study in rats. *Indian J Physiol Pharmacol* 29:234-238
 371. Gautam R, Pereira BM 1992 The effect of ovine prolactin on the epididymal sialic acid concentration in male rats. *Clin Exp Pharmacol Physiol* 19:495-501
 372. Gautam R, Pereira BM 1993 Modulation in activity of some epididymal glycosidases by prolactin. *Indian J Exp Biol* 31:410-413
 373. Ray B, Gautam R, Gaur M, Srivastava N, Pereira BM 1994 Impact of prolactin on epididymal lipid profile in castrated rats. *Indian J Exp Biol* 32:299-303
 374. Jayakumar PG, Arunakaran J, Aruldas MM, Govindarajulu P 1992 Role of prolactin on epididymal glycoprotein metabolism in matured monkeys, *Macaca radiata*: specific activities of glycosyltransferases and glycosidases. *Indian J Exp Biol* 30:1075-1078
 375. Segal S, Ron M, Laufer N, Ben-David M 1978 Prolactin in seminal plasma of infertile men. *Arch Androl* 1:49-52
 376. Sueldo CE, Berger T, Kletzky O, Marrs RP 1985 Seminal prolactin concentration and sperm reproductive capacity. *Fertil Steril* 43:632-635
 377. Chan SY, Tang LC, Chan PH, Tang GW, Ma HK 1984 Relationships of seminal plasma prolactin with spermatozoal characteristics and fertilizing capacity *in vitro*. *Arch Androl* 12:17-24
 378. Arunakaran J, Balasubramanian K, Srinivasan N, Aruldas MM, Govindarajulu P 1989 Effects of prolactin and androgens on seminal vesicular lipids of castrated mature bonnet monkeys *Macaca radiata*. *Indian J Exp Biol* 27:329-333
 379. Arunakaran J, Balasubramanian K, Srinivasan N, Aruldas MM, Govindarajulu P 1988 Effects of androgens, prolactin and bromocriptine on seminal vesicular enzymes of the pyruvate malate cycle involved in lipogenesis in castrated mature monkeys, *Macaca radiata*. *Int J Androl* 11:133-139
 380. Arunakaran J, Aruldas MM, Govindarajulu P 1988 Interactions of androgens and prolactin in the seminal vesicles of mature bonnet monkeys, *Macaca radiata*. I. Nucleic acids and phosphatases. *J Androl* 9:121-125
 381. Tam CC, Wong YC, Tang F 1992 Ultrastructural and cytochemical studies of the effects of prolactin on the lateral prostate and the seminal vesicle of the castrated guinea pig. *Cell Tissue Res* 270:105-112
 382. Resnick MI, Walvoord DJ, Grayhack JT 1974 Effect of prolactin on testosterone uptake by the perfused canine prostate. *Surg Forum* 25:70-72
 383. Johansson R 1976 Effect of prolactin, growth hormone and insulin on the uptake and binding of dihydrotestosterone to the cultured rat ventral prostate. *Acta Endocrinol (Copenh)* 81:854-864
 384. Prins GS, Lee C 1982 Influence of prolactin-producing pituitary grafts on the *in vivo* uptake, distribution, and disappearance of [³H]testosterone and [³H]dihydrotestosterone by the rat prostate lobes. *Endocrinology* 110:920-925
 385. Prins GS 1987 Prolactin influence on cytosol and nuclear androgen receptors in the ventral, dorsal, and lateral lobes of the rat prostate. *Endocrinology* 120:1457-1464
 386. Reiter E, Bonnet P, Sente B, Dombrowicz D, de Leval J, Closset J, Hennen G 1992 Growth hormone and prolactin stimulate androgen receptor, insulin-like growth factor-I (IGF-I) and IGF-I receptor levels in the prostate of immature rats. *Mol Cell Endocrinol* 88:77-87
 387. Tangbanluekal L, Robinette CL 1993 Prolactin mediates estradiol-induced inflammation in the lateral prostate of Wistar rats. *Endocrinology* 132:2407-2416
 388. Edwards WD, Thomas JA 1980 Morphologic and metabolic characteristics of ventral, lateral, dorsal and anterior prostate transplants in rats. Effect of testosterone and/or prolactin. *Horm Res* 13:28-39
 389. Dhananjaya Reddy Y, Venkatarami Reddy K, Govindappa S 1984 Metabolic profiles of prostate gland after the administration of prolactin and bromocriptine. *Arch Int Physiol Biochim* 92:125-130
 390. Srinivasan N, Aruldas MM, Govindarajulu P 1987 Interaction of sex steroids and prolactin on phosphatases, transaminases, and

- citric acid in the ventral prostate of male albino rats. Prostate 11:23-31
391. Costello LC, Franklin RB 1994 Effect of prolactin on the prostate. Prostate 24:162-166
 392. Franklin RB, Costello LC 1990 Prolactin directly stimulates citrate production and mitochondrial aspartate aminotransferase of prostate epithelial cells. Prostate 17:13-18
 393. Liu Y, Costello LC, Franklin RB 1996 Prolactin specifically regulates citrate oxidation and m-aconitase of rat prostate epithelial cells. Metabolism 45:442-449
 394. Costello LC, Liu Y, Franklin RB 1995 Prolactin specifically increases pyruvate dehydrogenase E1 α in rat lateral prostate epithelial cells. Prostate 26:189-193
 395. Reiter E, Lardinois S, Klug M, Sente B, Hennuy B, Bruyninx M, Closset J, Hennen G 1995 Androgen-independent effects of prolactin on the different lobes of the immature rat prostate. Mol Cell Endocrinol 112:113-122
 396. Vollmer EP, Gordon AS, Charipper HA 1942 Effects of hormones on erythropoiesis in the hypophysectomized rat. Endocrinology 31:619-628
 397. Dardenne M, Savino W, Gagnerault MC, Itoh T, Bach JF 1989 Neuroendocrine control of thymic hormonal production. I: Prolactin stimulates *in vivo* and *in vitro* the production of thymulin by human and murine thymic epithelial cells. Endocrinology 125:3-12
 398. Nagy E, Berczi I, Sabbadini E 1992 Endocrine control of the immunosuppressive activity of the submandibular gland. Brain Behav Immun 6:418-428
 399. Berczi I, Nagy E 1982 A possible role of prolactin in adjuvant arthritis. Arthritis Rheum 25:591-594
 400. Parkes D 1979 Drug therapy: bromocriptine. N Engl J Med 301:873-878
 401. Berczi I, Nagy E, Asa SL, Kovacs K 1984 The influence of pituitary hormones in adjuvant arthritis. Arthritis Rheum 27:682-688
 402. Nagy E, Berczi I 1981 Prolactin and contact sensitivity. Allergy 36:429-431
 403. Nagy E, Berczi I, Friesen HG 1983 Regulation of immunity in rats by lactogenic and growth hormones. Acta Endocrinol (Copenh) 102:351-357
 404. Berczi I, Nagy E, Kovacs K, Horvath E 1981 Regulation of humoral immunity in rats by pituitary hormones. Acta Endocrinol (Copenh) 98:506-513
 405. McMurray R, Keisler D, Kanuckel K, Izui S, Walker SE 1991 Prolactin influences autoimmune disease activity in the female B/W mouse. J Immunol 147:3780-3787
 406. Lahat N, Miller A, Shiller R, Touby E 1993 Differential effects of prolactin upon activation and differentiation of human B lymphocytes. J Neuroimmunol 47:35-40
 407. Nagy E, Berczi I 1989 Pituitary dependence of bone marrow function. Br J Haematol 71:457-462
 408. Bernton EW, Meltzer MS, Holaday JW 1988 Suppression of macrophage activation and T-lymphocyte function in hypoprolactinemic mice. Science 239:401-404
 409. Sabharwal P, Glaser R, LaFuse W, Varma S, Liu Q, Arkins S, Kooijman R, Kutz L, Kelley KW, Malarkey WB 1992 Prolactin synthesized and secreted by human peripheral blood mononuclear cells: an autocrine growth factor for lymphoproliferation. Proc Natl Acad Sci USA 89:7713-7716
 410. Murphy WJ, Durum SK, Longo DL 1993 Differential effects of growth hormone and prolactin on murine T cell development and function. J Exp Med 178:231-236
 411. Gala RR, Shevach EM 1993 Influence of prolactin and growth hormone on the activation of dwarf mouse lymphocytes *in vivo*. Proc Soc Exp Biol Med 204:224-230
 412. Bellone G, Geuna M, Carbone A, Silvestri S, Foa R, Emanuelli G, Matera L 1995 Regulatory action of prolactin on the *in vitro* growth of CD34+ve human hemopoietic progenitor cells. J Cell Physiol 163:221-231
 413. Gagnerault MC, Touraine P, Savino W, Kelly PA, Dardenne M 1993 Expression of prolactin receptors in murine lymphoid cells in normal and autoimmune situations. J Immunol 150:5673-5681
 414. O'Neal KD, Schwarz LA, Yu-Lee L-Y 1991 Prolactin receptor gene expression in lymphoid cells. Mol Cell Endocrinol 82:127-135
 415. Fletcher-Chiappini SE, Compton MM, LaVoie HA, Day EB, Witorsch RJ 1993 Glucocorticoid-prolactin interactions in Nb2 lymphoma cells: Antiproliferative vs. anticytolytic effects. Proc Soc Exp Biol Med 202:345-352
 416. LaVoie HA, Witorsch RJ 1995 Investigation of intracellular signals mediating the anti-apoptotic action of prolactin in Nb2 lymphoma cells. Proc Soc Exp Biol Med 209:257-269
 417. Cesario TC, Yousefi S, Carandang G, Sadati N, Le J, Vaziri N 1994 Enhanced yields of γ -interferon in prolactin treated human peripheral blood mononuclear cells. Proc Soc Exp Biol Med 205:89-95
 418. Comsa J, Leonhardt H, Schwarz JA 1975 Influence of the thymus-corticotropin-growth hormone interaction on the rejection of skin allografts in the rat. Ann NY Acad Sci 249:387-401
 419. Taub DD, Tsarfaty G, Lloyd AR, Durum SK, Longo DL, Murphy WJ 1994 Growth hormone promotes human T cell adhesion and migration to both human and murine matrix proteins *in vitro* and directly promotes xenogeneic engraftment. J Clin Invest 94:293-300
 420. Skwarlo-Sonta K 1990 Mitogenic effect of prolactin on chicken lymphocytes *in vitro*. Immunol Lett 24:171-177
 421. Witorsch RJ, Day EB, LaVoie HA, Hashemi N, Taylor JK 1993 Comparison of glucocorticoid-induced effects in prolactin-dependent and autonomous rat Nb2 lymphoma cells. Proc Soc Exp Biol Med 203:454-460
 422. Stevens AM, Wang YF, Sieger KA, Lu HF, Yu-Lee L-Y 1995 Biphasic transcriptional regulation of the interferon regulatory factor-1 gene by prolactin: involvement of γ -interferon-activated sequence and Stat-related proteins. Mol Endocrinol 9:513-525
 423. Stevens AM, Yu-Lee L-Y 1992 The transcription factor interferon regulatory factor-1 is expressed during both early G1 and the G1/S transition in the prolactin-induced lymphocyte cell cycle. Mol Endocrinol 6:2236-2243
 424. Andrews GK, Varma S, Ebner KE 1987 Regulation of expression of c-fos and c-myc in rat lymphoma Nb-2 cells. Biochim Biophys Acta 909:231-236
 425. Richards JF, Beer CT, Bourgeault C, Chen K, Gout PW 1982 Biochemical response of lymphoma cells to mitogenic stimulation by prolactin. Mol Cell Endocrinol 26:41-49
 426. deToledo SM, Murphy LJ, Hatton TH, Friesen HG 1987 Regulation of 70-kilodalton heat-shock-like messenger ribonucleic acid *in vitro* and *in vivo* by prolactin. Mol Endocrinol 1:430-434
 427. Yu-Lee L-Y 1988 Prolactin: role in T-cell proliferation. Ann NY Acad Sci 546:245-247
 428. Buckley AR, Buckley DJ, Leff MA, Hoover DS, Magnuson NS 1995 Rapid induction of *pim-1* expression by prolactin and interleukin-2 in rat Nb2 lymphoma cells. Endocrinology 136:5252-5259
 429. Gilks CB, Porter SD, Barker C, Tschlis PN, Gout PW 1995 Prolactin (PRL)-dependent expression of a zinc finger protein-encoding gene, Gfi-1, in Nb2 lymphoma cells: constitutive expression in autonomous sublines. Endocrinology 136:1805-1808
 430. Leff MA, Buckley DJ, Krumenacker JS, Reed JC, Miyashita T, Buckley AR 1996 Rapid modulation of the apoptosis regulatory genes, bcl-2 and bax by prolactin in rat Nb2 lymphoma cells. Endocrinology 137:5456-5462
 431. Hosokawa Y, Yang M, Kaneko S, Tanaka M, Nakashima K 1996 Prolactin induces switching of T-cell receptor gene expression from α to γ in rat Nb2 pre-T lymphoma cells. Biochem Biophys Res Commun 220:958-962
 432. Hosokawa Y, Onga T, Nakashima K 1994 Induction of D2 and D3 cyclin-encoding genes during promotion of the G1/S transition by prolactin in rat Nb2 cells. Gene 147:249-252
 433. Hosokawa Y, Yang M, Kaneko S, Tanaka M, Nakashima K 1995 Synergistic gene expressions of cyclin E, cdk2, cdk5 and E2F-1 during the prolactin-induced G1/S transition in rat Nb2 pre-T lymphoma cells. Biochem Mol Biol Int 37:393-399
 434. Axtell SM, Truong TM, O'Neal KD, Yu-Lee L-Y 1995 Characterization of a prolactin-inducible gene, clone 15, in T cells. Mol Endocrinol 9:312-318
 435. Wilson TM, Yu-Lee L-Y, Kelley MR 1995 Coordinate gene expression of luteinizing hormone-releasing hormone (LHRH) and the LHRH-receptor after prolactin stimulation in the rat Nb2 T-cell line: implications for a role in immunomodulation and cell cycle gene expression. Mol Endocrinol 9:44-53
 436. David M, Petricoin III EF, Igarashi K, Feldman GM, Finbloom DS, Larner AC 1994 Prolactin activates the interferon-regulated

- p91 transcription factor and the Jak2 kinase by tyrosine phosphorylation. *Proc Natl Acad Sci USA* 91:7174-7178
437. DaSilva L, Howard OM, Rui H, Kirken RA, Farrar WL 1994 Growth signaling and Jak2 association mediated by membrane-proximal regions of prolactin receptors. *J Biol Chem* 269:18267-18270
 438. Campbell GS, Argetsinger LS, Ihle JN, Kelly PA, Rillema JA, Carter-Su C 1994 Activation of JAK2 tyrosine kinase by prolactin receptors in Nb2 cells and mouse mammary gland explants. *Proc Natl Acad Sci USA* 91:5232-5236
 439. Lebrun JJ, Ali S, Sofer L, Ullrich A, Kelly PA 1994 Prolactin-induced proliferation of Nb2 cells involves tyrosine phosphorylation of the prolactin receptor and its associated tyrosine kinase. *J Biol Chem* 269:14021-14026
 440. Clevenger CV, Medaglia MV 1994 The protein tyrosine kinase p59fyn is associated with prolactin (PRL) receptor and is activated by PRL stimulation of T-lymphocytes. *Mol Endocrinol* 8:674-681
 441. Wang Y-F, Yu-Lee L-Y 1996 Multiple Stat complexes interact at the interferon regulatory factor-1 interferon- γ activation sequence in prolactin-stimulated Nb2 T cells. *Mol Cell Endocrinol* 121:19-28
 442. Gouilleux F, Moritz D, Humar M, Moriggi R, Berchtold S, Groner B 1995 Prolactin and interleukin-2 receptors in T lymphocytes signal through a MGF-STAT5-like transcription factor. *Endocrinology* 136:5700-5708
 443. Rao Y-P, Buckley DJ, Buckley AR 1995 Rapid activation of mitogen-activated protein kinase and p21^{ras} by prolactin and interleukin-2 in rat Nb2 node lymphoma cells. *Cell Growth Differ* 6:1235-1244
 444. Erwin RA, Kirken RA, Malabarba MG, Farrar WL, Rui H 1995 Prolactin activates Ras via signaling proteins SHC, growth factor receptor bound 2, and son of sevenless. *Endocrinology* 136:3512-3518
 445. Clevenger CV, Torigoe T, Reed JC 1994 Prolactin induces rapid phosphorylation and activation of prolactin receptor-associated RAF-1 kinase in a T-cell line. *J Biol Chem* 269:5559-5565
 446. Clevenger CV, Ngo W, Sokol DL, Luger SM, Gewirtz AM 1995 Vav is necessary for prolactin-stimulated proliferation and is translocated into the nucleus of a T-cell line. *J Biol Chem* 270:13246-13253
 447. Buckley AR, Rao Y-P, Buckley DJ, Gout PW 1994 Prolactin-induced phosphorylation and nuclear translocation of MAP kinase in Nb2 lymphoma cells. *Biochem Biophys Res Commun* 204:1158-1164
 448. Puissant C, Mitev V, Lemnaouar M, Manceau V, Sobel A, Houdebine LM 1995 Stathmin gene expression in mammary gland and in Nb2 cells. *Biol Cell* 85:109-115
 449. Meyer N, Prentice DA, Fox MT, Hughes JP 1992 Prolactin-induced proliferation of the Nb2 T-lymphoma is associated with protein kinase-C-independent phosphorylation of stathmin. *Endocrinology* 131:1977-1984
 450. Berlanga JJ, Gualillo O, Buteau H, Applanat M, Kelly PA, Edery M 1997 Prolactin activates tyrosyl phosphorylation of insulin receptor substrate-1 and phosphatidylinositol-3-OH kinase. *J Biol Chem* 272:2050-2052
 451. Ali S, Chen Z, Lebrun JJ, Vogel W, Kharitonov A, Kelly PA, Ullrich A 1996 PTP1D is a positive regulator of the prolactin signal leading to β -casein promoter activation. *EMBO J* 15:135-142
 452. Ganguli S, Hu L, Menke P, Collier RJ, Gertler A 1996 Nuclear accumulation of multiple protein kinases during prolactin-induced proliferation of Nb2 rat lymphoma cells. *J Cell Physiol* 167:251-260
 453. Buckley AR, Putnam CW, Russell DH 1988 Prolactin as a mammalian mitogen and tumor promoter. *Adv Enzyme Regul* 27:371-391
 454. Mitev V, Bayat-Sarmadi M, Lemnaouar M, Puissant C, Houdebine LM 1996 The effect of prolactin on casein kinase II, MAP kinase and PKC in rabbit mammary cells and Nb2 rat lymphoid cells. *Biochem Pharmacol* 52:1719-1727
 455. Al-Sakkaf KA, Dobson PRM, Brown BL 1996 Activation of phosphatidylinositol 3-kinase by prolactin in Nb2 cells. *Biochem Biophys Res Commun* 221:779-784
 456. Hunter S, Koch BL, Anderson SM 1997 Phosphorylation of cbl after stimulation of Nb2 cells with prolactin and its association with phosphatidylinositol 3-kinase. *Mol Endocrinol* 11:1213-1222
 457. Carey GB, Liberti JP 1995 Stimulation of receptor-associated kinase, tyrosine kinase, and MAP kinase is required for prolactin-mediated macromolecular biosynthesis and mitogenesis in Nb2 lymphoma. *Arch Biochem Biophys* 316:179-189
 458. Too CKL, Murphy PR, Friesen HG 1989 G-proteins modulate prolactin- and interleukin-2-stimulated mitogenesis in rat Nb2 lymphoma cells. *Endocrinology* 124:2185-2192
 459. Larsen JL, Dufau ML 1988 Modulation of prolactin-stimulated Nb2 lymphoma cell mitogenesis by cholera toxin and pertussis toxin. *Endocrinology* 123:438-444
 460. Hafez MM, Costlow ME 1989 Phosphatidylethanolamine turnover is an early event in the response of Nb2 lymphoma cells to prolactin. *Exp Cell Res* 184:37-43
 461. Too CKL, Cragoe Jr EJ, Friesen HG 1987 Amiloride-sensitive Na⁺/H⁺ exchange in rat Nb2 node lymphoma cells. Stimulation by prolactin and other mitogens. *Endocrinology* 121:1512-1520
 462. Oberholtzer E, Contarini M, Veglia F, Cossarizza A, Franceschi C, Geuna M, Provinciali M, Di Stefano G, Sissom J, Brizzi MF, Pegoraro L, Matera L 1996 Prolactin increases the susceptibility of primary leukemia cells to NK and LAK effectors. *Adv Neuroimmunol* 6:233-247
 463. Cesano A, Oberholtzer E, Contarini M, Geuna M, Bellone G, Matera L 1994 Independent and synergistic effect of interleukin-2 and prolactin on development of T- and NK-derived LAK effectors. *Immunopharmacology* 28:67-75
 464. Matera L, Cesano A, Bellone G, Oberholtzer E 1992 Modulatory effect of prolactin on the resting and mitogen-induced activity of T, B, and NK lymphocytes. *Brain Behav Immun* 6:409-417
 465. Sedo A, Van Weyenbergh J, Rouillard D, Bauvois B 1996 Synergistic effect of prolactin on IFN- γ -mediated growth arrest in human monoblastic cells: correlation with the up-regulation of IFN- γ receptor gene expression. *Immunol Lett* 53:125-130
 466. Edwards III CK, Ghiasuddin SM, Schepper JM, Yunger LM, Kelley KW 1988 A newly defined property of somatotropin: priming of macrophages for production of superoxide anion. *Science* 239:769-771
 467. Fu YK, Arkins S, Fuh G, Cunningham BC, Wells JA, Fong S, Cronin MJ, Dantzer R, Kelley KW 1992 Growth hormone augments superoxide anion secretion of human neutrophils by binding to the prolactin receptor. *J Clin Invest* 89:451-457
 468. Meli R, Raso GM, Bentivoglio C, Nuzzo I, Galdiero M, Di Carlo R 1996 Recombinant human prolactin induces protection against *Salmonella typhimurium* infection in the mouse: role of nitric oxide. *Immunopharmacology* 34:1-7
 469. Cecilia Fornari M, Fernanda Palacios M, Diez RA, Intebi AD 1994 Decreased chemotaxis of neutrophils in acromegaly and hyperprolactinemia. *Eur J Endocrinol* 130:463-468
 470. Villa-Verde DM, Mello-Coelho V, Lagrota-Candido JM, Chammass R, Savino W 1995 The thymic nurse cell complex: an *in vitro* model for extracellular matrix-mediated intrathymic T cell migration. *Braz J Med Biol Res* 28:907-912
 471. Weisz-Carrington P, Roux ME, McWilliams M, Phillips-Quagliata JM, Lamm ME 1978 Hormonal induction of the secretory immune system in the mammary gland. *Proc Natl Acad Sci USA* 75:2928-2932
 472. Gordon EM, Johnson TR, Ramos LP, Schmeidler-Sapiro KT 1991 Enhanced expression of factor XII (Hageman factor) in isolated livers of estrogen- and prolactin-treated rats. *J Lab Clin Med* 117:353-358
 473. Zellweger R, Wichmann MW, Ayala A, DeMaso CM, Chaudry IH 1996 Prolactin: a novel and safe immunomodulating hormone for the treatment of immunodepression following severe hemorrhage. *J Surg Res* 63:53-58
 474. El-Garf A, Salah S, Shaarawy M, Zaki S, Anwer S 1996 Prolactin hormone in juvenile systemic lupus erythematosus: a possible relationship to disease activity and CNS manifestations. *J Rheumatol* 23:374-377
 475. Walker SE, Allen SH, McMurray RW 1993 Prolactin and autoimmune disease. *Trends Endocrinol Metab* 4:147-151
 476. Riskind PN, Massacesi L, Doolittle TH, Hauser SL 1991 The role of prolactin in autoimmune demyelination: suppression of experimental allergic encephalomyelitis by bromocriptine. *Ann Neurol* 29:542-547

477. McMurray RW, Allen SH, Pepmueller PH, Keisler D, Cassidy JT 1995 Elevated serum prolactin levels in children with juvenile rheumatoid arthritis and antinuclear antibody seropositivity. *J Rheumatol* 22:1577-1580
478. Neidhart M, Fluckiger EW 1992 Hyperprolactinaemia in hypophysectomized or intact male rats and the development of adjuvant arthritis. *Immunology* 77:449-455
479. Hinterberger-Fisher M, Kier P, Spona J, Sebesta C, Tiefengraber E, Habertheuer KH, Ruckser R, Schmid A, Kalhs P, Lechner K, Hinterberger W 1994 Prolactin: a possible mediator of graft-vs.-host disease following allogeneic bone marrow transplantation in humans. *Bone Marrow Transplant* 14:403-406
480. Gala RR, Shevach EM 1997 The immunomodulatory effects of a recombinant bovine prolactin (rbPRL) are secondary to endotoxin contamination. *FASEB J* 11:99A
481. Carrier M, Emery RW, Wild-Mobley J, Perrotta NJ, Russell DH, Copeland JG 1987 Prolactin as a marker of rejection in human heart transplantation. *Transplant Proc* 19:3442-3443
482. Yu-Lee L-Y 1997 Molecular actions of prolactin in the immune system. *Proc Soc Exp Biol Med* 215:35-51
483. Tanaka T, Shiu RP, Gout PW, Beer CT, Noble RL, Friesen HG 1980 A new sensitive and specific bioassay for lactogenic hormones: measurement of prolactin and growth hormone in human serum. *J Clin Endocrinol Metab* 51:1058-1063
484. Clevenger CV, Thickman K, Ngo W, Chang W-P, Takayama S, Reed JC 1997 Role of Bag-1 in the survival and proliferation of the cytokine-dependent lymphocyte lines, Ba/F3 and Nb2. *Mol Endocrinol* 11:608-618
485. Blackwell RE 1992 Hyperprolactinemia: evaluation and management. *Endocrinol Metab Clin North Am* 21:105-124
486. Healy DL, Herington AC, O'Herlihy C 1985 Chronic polyhydramnios is a syndrome with a lactogen receptor defect in the chorion laeve. *Br J Obstet Gynecol* 92:461-467
487. Nagano M, Kelly PA 1992 Absence of a putative ATP/GTP binding site in the rat prolactin receptor. *Biochem Biophys Res Commun* 183:610-618
488. Rui H, Djeu JY, Evans GA, Kelly PA, Farrar WL 1992 Prolactin receptor triggering: evidence for rapid tyrosine kinase activation. *J Biol Chem* 267:24076-24081
489. Carey GB, Liberti JP 1993 Inhibition of tyrosine kinase activity abolishes anabolic and mitogenic effects of prolactin and growth hormone in Nb2 cells. *Biochem Mol Biol Int* 31:635-642
490. Wilks AF, Harpur AG, Kurban RR, Ralph SJ, Zürcher G, Ziemiecki A 1991 Two novel protein-tyrosine kinases, each with a second phosphotransferase-related catalytic domain, define a new class of protein kinase. *Mol Cell Biol* 11:2057-2065
491. Harpur AG, Andres A-C, Ziemiecki A, Aston RR, Wilks AF 1992 Jak2, a third member of the Jak family of protein tyrosine kinases. *Oncogene* 7:1347-1353
492. Ihle JN, Witthuhn BA, Quelle FW, Yamamoto K, Thierfelder WE, Kreider B, Silvennoinen O 1994 Signalling by the cytokine receptor superfamily: Jaks and Stats. *Trends Biochem Sci* 19:222-227
493. Ihle JN 1994 Signaling by the cytokine receptor superfamily. Just another kinase story. *Trends Endocrinol Metab* 5:137-143
494. Kishimoto T, Taga T, Akira S 1994 Cytokine signal transduction. *Cell* 76:253-262
495. Ihle JN, Kerr IM 1995 Jaks and Stats in signaling by the cytokine receptor superfamily. *Trends Genet* 11:69-74
496. Argetsinger LS, Campbell GS, Yang X, Witthuhn BA, Silvennoinen O, Ihle JN, Carter-Su C 1993 Identification of Jak2 as a growth hormone receptor-associated tyrosine kinase. *Cell* 74:237-244
497. Rui H, Kirken RA, Farrar WL 1994 Activation of receptor-associated tyrosine kinase JAK2 by prolactin. *J Biol Chem* 269:5364-5368
498. Dusanter-Fourt I, Muller O, Ziemiecki A, Mayeux P, Drucker B, Djiane J, Wilks A, Harper AG, Fischer S, Gisselbrecht S 1994 Identification of Jak protein tyrosine kinases as signaling molecules for prolactin. Functional analysis of prolactin receptor and prolactin-erythropoietin receptor chimera expressed in lymphoid cells. *EMBO J* 13:2583-2591
499. Han Y, Watling D, Rogers NC, Stark GR 1997 Jak2 and Stat5, but not Jak1 and Stat1, are required for prolactin-induced β -lactoglobulin transcription. *Mol Endocrinol* 11:1180-1188
500. Goupille O, Daniel N, Bignon C, Jolivet G, Djiane J 1997 Prolactin signal transduction to milk protein genes: carboxy-terminal part of the prolactin receptor and its tyrosine phosphorylation are not obligatory for Jak2 and Stat5 activation. *Mol Cell Endocrinol* 127:155-169
501. Lebrun JJ, Ali S, Ullrich A, Kelly PA 1995 Proline-rich sequence-mediated JAK2 association to the prolactin receptor is required but not sufficient for signal transduction. *J Biol Chem* 270:10664-10670
502. Ren R, Mayer BJ, Cicchetti P, Baltimore D 1993 Identification of ten amino acid proline-rich SH3 binding site. *Science* 259:1157-1161
503. Pezet A, Buteau H, Kelly PA, Edery M 1997 The last proline of Box 1 is essential for association with JAK2 and functional activation of the prolactin receptor. *Mol Cell Endocrinol* 129:199-208
504. O'Neal KD, Chari MV, McDonald CH, Cook RG, Yu-Lee L-Y, Morrisett JD, Shearer WT 1996 Multiple *cis-trans* conformers of the prolactin receptor proline-rich motif (PRM) peptide detected by reverse-phase HPLC, CD, and NMR. *Biochem J* 315:833-844
505. Wang YD, Wood WI 1995 Amino acids of the human growth hormone receptor that are required for proliferation and Jak-STAT signaling. *Mol Endocrinol* 9:303-311
506. Dinerstein H, Lago F, Goujon L, Ferrag F, Esposito N, Finidori J, Kelly PA, Postel-Vinay MC 1995 The proline-rich region of the growth hormone receptor is essential for Jak2 phosphorylation, activation of cell proliferation and gene transcription. *Mol Endocrinol* 9:1701-1707
507. Wang Y-F, O'Neal KD, Yu-Lee L-Y 1997 Multiple prolactin receptor cytoplasmic residues and Stat1 mediate prolactin signaling to the IRF-1 promoter. *Mol Endocrinol* 11:1353-1364
508. DaSilva L, Rui H, Erwin RA, Zack Howard OM, Kirken RA, Malabarba MG, Hackett RH, Lerner AC, Farrar WL 1996 Prolactin recruits STAT1, STAT3 and STAT5 independent of conserved receptor tyrosines TYR402, TYR479, TYR515 and TYR580. *Mol Cell Endocrinol* 117:131-140
509. Frank SJ, Yi W, Zhao Y, Goldsmith JF, Gilliland G, Jiang J, Sakai I, Kraft AS 1995 Regions of the JAK2 tyrosine kinase required for coupling to the growth hormone receptor. *J Biol Chem* 270:14776-14785
- 509a. Ferrag F, Pezet A, Chiarenza A, Buteau H, Nelson BH, Goffin V, Kelly PA 1998 Homodimerization of the IL-2 receptor β chain is necessary and sufficient to activate JAK2 and downstream signaling pathways. *FEBS Lett* 421:32-36
510. Chang WP, Clevenger CV 1996 Modulation of growth factor receptor function by isoform heterodimerization. *Proc Natl Acad Sci USA* 93:5947-5952
- 510a. Chang WP, Ye Y, Clevenger CV 1998 Stoichiometric structure-function analysis of the prolactin receptor signaling domain by receptor chimeras. *Mol Cell Biol* 18:896-905
511. Perrot-Applanat M, Gualillo O, Pezet A, Vincent V, Edery M, Kelly PA 1997 Dominant negative and cooperative effects of mutant forms of prolactin receptor. *Mol Endocrinol* 11:1020-1032
512. Berlanga JJ, Garcia-Ruiz JP, Perrot-Applanat M, Kelly PA, Edery M 1997 The short form of the prolactin receptor silences prolactin induction of the β -casein gene promoter. *Mol Endocrinol* 11:1449-1457
513. Lebrun JJ, Ali S, Goffin V, Ullrich A, Kelly PA 1995 A single phosphotyrosine residue of the prolactin receptor is responsible for activation of gene transcription. *Proc Natl Acad Sci USA* 92:4031-4035
514. Pezet A, Ferrag F, Kelly PA, Edery M 1997 Tyrosine docking sites of the rat prolactin receptor required for association and activation of Stat5. *J Biol Chem* 272:25043-25050
515. Sotiropoulos A, Perrot-Applanat M, Dinerstein H, Pallier A, Postel-Vinay MC, Finidori J, Kelly PA 1994 Distinct cytoplasmic regions of the growth hormone receptor are required for activation of Jak2, mitogen-activated protein kinase, and transcription. *Endocrinology* 135:1292-1298
516. Sotiropoulos A, Moutoussamy S, Renaudie F, Clauss M, Kayser C, Guilleux F, Kelly PA, Finidori J 1996 Differential activation of Stat3 and Stat5 by distinct regions of the growth hormone receptor. *Mol Endocrinol* 10:998-1009
517. Lobie PE, Allevato G, Nielsen JH, Norstedt G, Billestrup N 1995 Requirement of tyrosine residues 333 and 338 of the growth hor-

- mone (GH) receptor for selected GH-stimulated function. *J Biol Chem* 270:21745-21750
518. Hackett RH, Wang YD, Larner AC 1995 Mapping of the cytoplasmic domain of the human growth hormone receptor required for the activation of JAK2 and Stat proteins. *J Biol Chem* 270:21326-21330
 519. VanderKuur JA, Wang X, Zhang L, Allevato G, Billestrup N, Carter-Su C 1995 Growth hormone-dependent phosphorylation of tyrosine 333 and/or 338 of the growth hormone receptor. *J Biol Chem* 270:21738-21744
 520. Smit LS, Meyer DJ, Billestrup N, Norstedt G, Schwartz J, Carter-Su C 1996 The role of the growth hormone (GH) receptor and Jak1 and Jak2 kinases in the activation of Stats 1, 3, and 5 by GH. *Mol Endocrinol* 10:519-533
 521. Wang X, Darus CJ, Xu BC, Kopchick JJ 1996 Identification of growth hormone receptor (GHR) tyrosine residues required for GHR phosphorylation and JAK2 and Stat5 activation. *Mol Endocrinol* 10:1249-1260
 522. Ali S, Edery M, Pellegrini I, Lesueur L, Paly J, Djiane J, Kelly PA 1992 The Nb2 form of prolactin receptor is able to activate a milk protein gene promoter. *Mol Endocrinol* 6:1242-1248
 523. Scott P, Kessler MA, Schuler LA 1992 Molecular cloning of the bovine prolactin receptor and distribution of prolactin and growth hormone receptor transcripts in fetal and utero-placental tissues. *Mol Cell Endocrinol* 89:47-58
 524. Jabbour HN, Clarke LA, Boddy S, Pezet A, Edery M, Kelly PA 1996 Cloning, sequencing and functional analysis of a truncated cDNA encoding red deer prolactin receptor: an alternative tyrosine residue mediates β -casein promoter activation. *Mol Cell Endocrinol* 123:17-26
 525. Das R, Vonderhaar BK 1995 Transduction of prolactin's (PRL) growth signal through both long and short forms of the PRL receptor. *Mol Endocrinol* 9:1750-1759
 526. O'Neal KD, Yu-Lee L-Y 1994 Differential signal transduction of the short, Nb2 and long prolactin receptors. *J Biol Chem* 269:26076-26082
 527. Yu-Lee L-Y, Hrachovy JA, Stevens A, Schwarz LA 1990 Interferon-regulatory factor 1 is an immediate-early gene under transcriptional regulation by prolactin in Nb2 T cells. *Mol Cell Biol* 10:3087-3094
 528. Gouilleux F, Wakao H, Mundt M, Groner B 1994 Prolactin induces phosphorylation of Tyr694 of Stat5(MGF), a prerequisite for DNA binding and induction of transcription. *EMBO J* 13:4361-4369
 529. Shuai K, Horvath CM, Tsai Huang LH, Qureshi SA, Cowburn D, Darnell Jr JE 1994 Interferon activation of the transcription factor Stat91 involves dimerization through SH2-phosphotyrosyl peptide interactions. *Cell* 76:821-828
 530. Darnell Jr JE, Kerr IM, Stark GR 1994 Jak-STAT pathways and transcriptional activation in response to IFNs and other extracellular signaling proteins. *Science* 264:1415-1421
 531. Taniguchi T 1995 Cytokine signaling through nonreceptor protein tyrosine kinases. *Science* 268:251-255
 532. Ihle JN 1996 STATs: Signal Transducers and Activators of Transcription. *Cell* 84:331-334
 533. Wakao H, Gouilleux F, Groner B 1994 Mammary gland factor (MGF) is a novel member of the cytokine regulated transcription factor gene family and confers the prolactin response. *EMBO J* 13:2182-2191
 534. Liu X, Robinson GW, Gouilleux F, Groner B, Hennighausen L 1995 Cloning and expression of Stat5 and an additional homologue (Stat5b) involved in prolactin signal transduction in mouse mammary tissue. *Proc Natl Acad Sci USA* 92:8831-8835
 535. Azam M, Erdjument-Bromage H, Kreider BL, Xia M, Quelle F, Basu R, Saris C, Tempst P, Ihle JN, Schindler C 1995 Interleukin-3 signals through multiple isoforms of Stat5. *EMBO J* 14:1402-1411
 536. Mui AL-F, Wakao H, O'Farrell A-M, Harada N, Miyajima A 1995 Interleukin-3, granulocyte-macrophage colony stimulating factor and interleukin-5 transduce signals through two STAT5 homologs. *EMBO J* 14:1166-1175
 537. Kazansky AV, Raught B, Lindsey SM, Wang YF, Rosen JM 1996 Regulation of mammary gland factor/Stat5a during mammary gland development. *Mol Endocrinol* 9:1598-1609
 538. Hou J, Schindler U, Henzel WJ, Wong SC, McKnight SL 1995 Identification and purification of human Stat proteins activated in response to interleukin-2. *Immunity* 2:321-329
 539. Silva CM, Lu H, Day RN 1996 Characterization and cloning of Stat5 from IM-9 cells and its activation by growth hormone. *Mol Endocrinol* 10:508-518
 540. Mui AL, Wakao H, Kinoshita T, Kitamura T, Miyajima A 1996 Suppression of interleukin-3-induced gene expression by a C-terminal truncated Stat5: role of Stat5 in proliferation. *EMBO J* 15:2425-2433
 541. Moriggi R, Gouilleux-Gruart V, Jähne R, Berchtold S, Gartmann C, Liu X, Hennighausen L, Sotiropoulos A, Groner B, Gouilleux F 1996 Deletion of the carboxyl-terminal transactivation domain of MGF-Stat5 results in sustained DNA binding and a dominant negative phenotype. *Mol Cell Biol* 16:5691-5700
 542. Wang D, Stravopodis D, Teglund S, Kitazawa J, Ihle JN 1996 Naturally occurring dominant negative variants of Stat5. *Mol Cell Biol* 16:6141-6148
 543. Ferrag F, Chiarenza A, Goffin V, Kelly PA 1996 Convergence of signaling transduced by PRL/cytokine chimeric receptors on PRL-responsive gene transcription. *Mol Endocrinol* 10:451-460
 544. Beadling C, Ng J, Babbage JW, Cantrell DA 1996 Interleukin-2 activation of STAT5 requires the convergent action of tyrosine kinases and a serine/threonine kinase pathway distinct from the Raf1/ERK2 MAP kinase pathway. *EMBO J* 15:1902-1913
 545. Wen Z, Zhong Z, Darnell Jr JE 1995 Maximal activation of transcription by Stat1 and Stat3 requires both tyrosine and serine phosphorylation. *Cell* 82:241-250
 546. Schmitt-Ney M, Happ B, Ball RK, Groner B 1992 Developmental and environmental regulation of a mammary gland-specific nuclear factor essential for transcription of the gene encoding β -casein. *Proc Natl Acad Sci USA* 89:3130-3134
 547. Marte BM, Meyer T, Stabel S, Standke GJ, Jaken S, Fabbro D, Hynes NE 1994 Protein kinase C and mammary cell differentiation: involvement of protein kinase C α in the induction of β -casein gene expression. *Cell Growth Differ* 5:239-247
 548. Quelle FW, Wang D, Nosaka T, Thierfelder WE, Stravopodis D, Weinstein Y, Ihle JN 1996 Erythropoietin induces activation of Stat5 through association with specific tyrosines on the receptor that are not required for a mitogenic response. *Mol Cell Biol* 16:1622-1631
 549. Liu X, Robinson GW, Hennighausen L 1996 Activation of Stat5a and Stat5b by tyrosine phosphorylation is tightly linked to mammary gland differentiation. *Mol Endocrinol* 10:1496-1506
 550. Kirken RA, Malabarba MG, Xu J, Liu X, Farrar WL, Hennighausen L, Larner AC, Grimley PM, Rui H 1997 Prolactin stimulates serine/tyrosine phosphorylation and formation of hetero-complexes of multiple Stat5 isoforms in Nb2 lymphocytes. *J Biol Chem* 272:14098-14103
 551. Luo G, Yu-Lee L-Y 1997 Transcriptional inhibition by Stat5: differential activities at growth-related vs. differentiation-specific promoters. *J Biol Chem* 272:26841-26849
 552. Lütticken C, Wegenka UM, Yuan J, Buschmann J, Schindler C, Ziemiecki Harpur AG, Wilks AF, Yasukawa K, Taga T, Kishimoto T, Barbieri G, Pellegrini S, Sendtner M, Heinrich PC, Horn F 1994 Association of transcription factor APRF and protein kinase Jak1 with the interleukin-6 signal transducer gp130. *Science* 263:89-92
 553. Levy DE, Kessler DS, Pine R, Darnell Jr JE 1989 Cytoplasmic activation of ISGF3, the positive regulator of interferon- α stimulated transcription, reconstituted *in vitro*. *Genes Dev* 3:1362-1371
 554. Gupta S, Yan H, Wong LH, Ralph S, Krolewski J, Schindler C 1996 The SH2 domains of Stat1 and Stat2 mediate multiple interactions in the transduction of INF- α signals. *EMBO J* 15:1075-1084
 555. Yi W, Kim SO, Jiang J, Park SH, Kraft AS, Waxman DJ, Frank SJ 1996 Growth hormone receptor cytoplasmic domain differentially promotes tyrosine phosphorylation of signal transducers and activators of transcription 5b and 3 by activated JAK2 kinase. *Mol Endocrinol* 10:1425-1443
 556. Songyang Z, Shoelson SE, Chaudhuri M, Gish G, Pawson T, Haser WG, King F, Roberts T, Ratnofsky S, Lechleider RJ, Neel BG, Birge RB, Fajardo JE, Chou MM, Hanafusa H, Schaffhausen B, Cantley LC 1993 SH2 domains recognize specific phosphopeptides sequences. *Cell* 72:767-778

557. Songyang Z, Shoelson SE, McGlade J, Olivier P, Pawson T, Bustelo XR, Barbacid M, Sabe H, Hanafusa H, Yi T, Ren R, Baltimore D, Ratnoffsky S, Feldman RA, Cantley LC 1994 Specific motifs recognized by the SH2 domains of Csk, 3BP2, fps/fes, GRB2, HCP, Syk, and Vav. *Mol Cell Biol* 14:2777-2785
558. Stahl N, Farruggella TJ, Boulton TG, Zhong Z, Darnell Jr JE, Yancopoulos GD 1995 Choice of STATs and other substrates specified by modular tyrosine-based motifs in cytokine receptors. *Science* 267:1349-1353
559. Gerhartz C, Heesel B, Sasse J, Hemmann U, Landgraf C, Schneider-Mergener J, Horn F, Heinrich PC, Graeve L 1996 Differential activation of acute phase response factor/STAT3 and STAT1 via the cytoplasmic domain of the interleukin 6 signal transducer gp130. I. Definition of a novel phosphotyrosine motif mediating Stat1 activation. *J Biol Chem* 271:12991-12998
560. Hemmann U, Gerhartz C, Heesel B, Sasse J, Kurapat G, Grötzinger J, Wollmer A, Zhong Z, Darnell Jr JE, Graeve L, Heinrich PC, Horn F 1996 Differential activation of acute phase response factor/STAT3 and STAT1 via the cytoplasmic domain of the interleukin 6 signal transducer gp130. II. Src homology SH2 domains define the specificity of STAT factor activation. *J Biol Chem* 271:12999-13007
561. May P, Gerhartz C, Heesel B, Welte T, Doppler W, Graeve L, Horn F, Heinrich PC 1996 Comparative study on the phosphotyrosine motifs of different cytokine receptors involved in Stat5 activation. *FEBS Lett* 394:221-226
562. Gobert S, Chretien S, Gouilleux F, Muller O, Pallard C, Dusanter-Fourt I, Groner B, Lacombe C, Gisselbrecht S, Mayeux P 1996 Identification of tyrosine residues within the intracellular domain of the erythropoietin receptor crucial for Stat5 activation. *EMBO J* 15:2434-2441
563. Damen JE, Wakao H, Miyajima A, Kros J, Humphries RK, Cutler RL, Krystal G 1995 Tyrosine 343 in the erythropoietin receptor positively regulates erythropoietin-induced cell proliferation and Stat5 activation. *EMBO J* 14:5557-5568
564. Fujii H, Nakagawa Y, Schindler U, Kawahara A, Mori H, Gouilleux F, Groner B, Ihle JN, Minami Y, Miyazaki T, Taniguchi T 1995 Activation of Stat 5 by interleukin 2 requires a carboxy-terminal region of the interleukin 2 receptor β chain but is not essential for the proliferative signal transmission. *Proc Natl Acad Sci USA* 92:5482-5486
565. Silvennoinen O, Witthuhn BA, Quelle FW, Cleveland JL, Yi T, Ihle JN 1993 Structure of the murine Jak2 protein-tyrosine kinase and its role in interleukin 3 signal transduction. *Proc Natl Acad Sci USA* 90:8429-8433
566. Horseman ND, Yu-Lee L-Y 1994 Transcriptional regulation by the helix bundle peptide hormones: growth hormone, prolactin, and hematopoietic cytokines. *Endocr Rev* 15:627-649
567. Gouilleux F, Pallard C, Dusanter-Fourt I, Wakao H, Haldosen L-A, Norstedt G, Levy D, Groner B 1995 Prolactin, growth hormone, erythropoietin and granulocyte-macrophage colony stimulating factor induce MGF-Stat5 DNA binding activity. *EMBO J* 14:2005-2013
568. Pallard C, Gouilleux F, Charon M, Groner B, Gisselbrecht S, Dusanter-Fourt I 1995 Interleukin-3, erythropoietin and prolactin activate a STAT5-like factor in lymphoid cells. *J Biol Chem* 270:15942-15945
569. Wakao H, Harada N, Kitamura T, Mui AL-F, Miyajima A 1995 Interleukin 2 and erythropoietin activate Stat5/MGF via distinct pathways. *EMBO J* 14:2527-2535
570. Wood TJJ, Sliva D, Lobie PE, Fircher TJ, Gouilleux F, Wakao H, Gustafsson JA, Groner B, Norstedt G, Haldosen L-A 1995 Mediation of growth hormone-dependent transcriptional activation by mammary gland factor/Stat5. *J Biol Chem* 270:9448-9453
571. Galsgaard ED, Gouilleux F, Groner B, Serup P, Nielsen JH, Billestrup N 1996 Identification of a growth hormone-responsive STAT5-binding element in the rat insulin 1 gene. *Mol Endocrinol* 10:652-660
572. Stöcklin E, Wissler M, Gouilleux F, Groner B 1996 Functional interactions between Stat5 and the glucocorticoid receptor. *Nature* 383:726-728
- 572a. Stöcklin E, Wissler M, Morrigl R, Groner B 1997 Specific DNA binding of Stat5, but not of the glucocorticoid receptor, is required for their functional cooperation in the regulation of gene transcription. *Mol Cell Biol* 17:6708-6716
573. Lechner J, Welte T, Tomasi JK, Bruno P, Cairns C, Gustafsson JA, Doppler W 1997 Promoter-dependent synergy between glucocorticoid receptor and Stat5 in the activation of β -casein gene transcription. *J Biol Chem* 272:20954-20960
574. Avruch J, Zhang X-F, Kyriakis JM 1994 Raf meets Ras: completing the framework of a signal transduction pathway. *Trends Biochem Sci* 19:279-283
575. Das R, Vonderhaar BK 1996 Activation of raf-1, MEK, and MAP kinase in prolactin responsive mammary cells. *Breast Cancer Res Treat* 40:141-149
576. Ihle JN 1996 STATs and MAPKs: obligate or opportunistic partners in signaling. *Bioessays* 18:95-98
577. Erpel T, Courtneidge SA 1995 Src family protein tyrosine kinases and cellular signal transduction pathways. *Curr Opin Cell Biol* 7:176-182
- 577a.al-Sakkaf KA, Dobson PR, Brown BL 1997 Prolactin tyrosine phosphorylation of p59fyn may mediate phosphatidylinositol 3-kinase activation in Nb2 cells. *J Mol Endocrinol* 19:347-350
578. Tourkine N, Schindler C, Larose M, Houdebine LM 1995 Activation of Stat factors by prolactin, interferon- γ , growth hormones, and a tyrosine phosphatase inhibitor in rabbit primary mammary epithelial cells. *J Biol Chem* 270:20952-20961
579. Daniel N, Waters MJ, Bignon C, Djiane J 1996 Involvement of a subset of tyrosine kinases and phosphatases in regulation of the β -lactoglobulin gene promoter by prolactin. *Mol Cell Endocrinol* 118:25-35
580. Adachi M, Fisher EH, Ihle JN, Imai K, Jirik F, Neel B, Pawson T, Shen SH, Thomas M, Ullrich A, Zhao Z 1996 Mammalian SH2-containing protein tyrosine phosphatases. *Cell* 85:15
- 580a. Yoshimura A, Ohkubo T, Kiguchi T, Jenkins NA, Gilbert DJ, Copeland NG, Hara T, Miyajima A 1995 A novel cytokine-inducible gene CIS encodes an SH2-containing protein that binds to tyrosine-phosphorylated interleukin 3 and erythropoietin receptors. *EMBO J* 14:2816-2826
- 580b. Starr R, Willson TA, Viney E, Murray LJJ, Rayner JR, Jenkins BJ, Gonda TJ, Alexander WS, Metcalf D, Nicola NA, Hilton DJ 1997 A family of cytokine-inducible inhibitors of signalling. *Nature* 387:917-921
- 580c. Endo TA, Masuhara M, Yokouchi M, Suzuki R, Sakamoto H, Mitsui K, Matsumoto A, Tanimura S, Ohtsubo M, Misawa H, Miyazaki T, Leonor N, Taniguchi T, Fujita T, Kanakura Y, Komiya S, Yoshimura A 1997 A new protein containing an SH2 domain that inhibits JAK kinases. *Nature* 387:921-924
- 580d. Naka T, Narazaki M, Hirata M, Matsumoto T, Minamoto S, Aono A, Nishimoto N, Kajita T, Taga T, Yoshizaki K, Akira S, Kishimoto T 1997 Structure and function of a new STAT-induced STAT inhibitor. *Nature* 387:924-927
- 580e. Chung CD, Liao J, Liu B, Jay P, Berta P, Shuai K 1997 Specific inhibition of Stat3 signal transduction by PIAS3. *Science* 248:1803-1805
581. Rao Y-P, Buckley DJ, Olson MD, Buckley AR 1995 Nuclear translocation of prolactin: collaboration of tyrosine kinase and protein kinase C activation in rat Nb2 node lymphoma cells. *J Cell Physiol* 163:266-276
582. Vacher P, Tran Van Chuoi M, Paly J, Djiane J, Dufy B 1994 Short term effect of prolactin on intracellular calcium in Chinese hamster ovary cells stably transfected with prolactin receptor complementary deoxyribonucleic acid. *Endocrinology* 134:1213-1218
583. Too CKL, Shiu RPC, Friesen HG 1990 Cross-linking of G-proteins to the prolactin receptor in rat Nb2 lymphoma cells. *Biochem Biophys Res Commun* 173:48-52
584. Too CKL, Murphy PR, Friesen HG 1990 The role of G-proteins in the mitogenesis of rat lactogen-dependent and lactogen-independent Nb2 lymphoma cells. *Endocrinology* 126:1368-1373
585. Duan WR, Linzer DIH, Gibori G 1996 Cloning and characterization of an ovarian-specific protein that associates with the short form of the prolactin receptor. *J Biol Chem* 271:15602-15607
- 585a. Quijano VJ, Sheffield LG 1998 Prolactin decreases epidermal growth factor receptor kinase activity via phosphorylation-dependent mechanism. *J Biol Chem* 273:1200-1207
- 585b. Yamauchi T, Ueki K, Tobe K, Tamemoto H, Sekine N, Wada M,

- Honjo M, Takahashi M, Takahashi T, Hirai H, Tushima T, Akanuma Y, Fujita T, Komuro I, Yazaki Y, Kadowaki T 1997 Tyrosine phosphorylation of the EGF receptor by the kinase Jak2 is induced by growth hormone. *Nature* 390:91-96
586. Nicoll CS 1980 Ontogeny and evolution of prolactin's functions. *Fed Proc* 39:2563-2566
587. Pellegrini I, Lebrun JJ, Ali S, Kelly PA 1992 Expression of prolactin and its receptor in human lymphoid cells. *Mol Endocrinol* 6:1023-1031
588. Ginsburg E, Vonderhaar BK 1995 Prolactin synthesis and secretion by human breast cancer cells. *Cancer Res* 55:2591-2595
589. Thomas K, Capocchi M 1987 Site-directed mutagenesis by gene targeting in mouse embryo-derived stem cells. *Cell* 51:503-512
590. Buck K, Vanek M, Groner B, Ball RK 1992 Multiple forms of prolactin receptor messenger ribonucleic acid are specifically expressed and regulated in murine tissues and the mammary cell line HC11. *Endocrinology* 130:1108-1114
591. Clarke DL, Arey BJ, Linzer DIH 1993 Prolactin receptor messenger ribonucleic acid expression in the ovary during rat estrous cycle. *Endocrinology* 133:2594-2603
592. Okamura H, Zachwieja J, Raguet S, Kelly PA 1989 Characterization and application of monoclonal antibodies to the prolactin receptor. *Endocrinology* 124:2499-2508
593. Nandi S 1958 Endocrine control of mammary gland development and function in the C³H/He Crgl mouse. *J Natl Cancer Inst* 21:1039-1063
594. Vonderhaar B 1988 Regulation of development of the normal mammary gland by hormones and growth factors. In: Lippman M, Dickson R (eds) *Breast Cancer: Cellular and Molecular Biology*. Kluwer Academic Publishers, Boston, MA, pp 252-266
595. Ormandy CJ, Sutherland RL 1993 Mechanisms of prolactin receptor regulation in mammary gland. *Mol Cell Endocrinol* 91:C1-C6
596. Lydon JP, DeMayo FJ, Funk CR, Mani SK, Hughes CA, Montgomery CA, Shyamala G, Conneely OM, O'Malley BW 1995 Mice lacking progesterone receptor exhibit pleiotropic reproductive abnormalities. *Genes Dev* 9:2266-2278
597. Gunnet JW, Freeman ME 1983 The mating-induced release of prolactin: a unique neuroendocrine response. *Endocr Rev* 4:44-61
598. Laufer N, Botero RW, DeCherney AH, Haseltine F, Polan ML, Behrman HR 1984 Gonadotropin and prolactin levels in follicular fluid of human ova successfully fertilized *in vitro*. *J Clin Endocrinol Metab* 58:430-434
599. Bohnet HG, Baukloh V, Kemeter P, Feichtinger W 1985 Prolactin and sex steroid concentrations in follicular fluids after different ovarian stimulation methods in IVF patients. *Ann NY Acad Sci* 442:227-235
600. Reinthaller A, Deutinger J, Bieglmayer C, Riss P, Muller TE, Fischl F, Janisch H 1987 Hormonal parameters in follicular fluid and the fertilization rate of *in vitro* cultured oocytes. *Arch Gynecol* 240:207-210
601. Reinthaller A, Deutinger J, Riss P, Muller TE, Fischl F, Bieglmayer C, Janisch H 1987 Relationship between the steroid and prolactin concentration in follicular fluid and the maturation and fertilization of human oocytes. *J In Vitro Fert Embryo Transfer* 4:228-231
602. Gonen Y, Casper RF 1989 Does transient hyperprolactinemia during ovarian hyperstimulation interfere with conception or pregnancy outcome? *Fertil Steril* 51:1007-1010
603. Seibel MM, Smith D, Dlugi AM, Levesque L 1989 Periovulatory follicular fluid hormone levels in spontaneous human cycles. *J Clin Endocrinol Metab* 68:1073-1077
604. Oda T, Yoshimura Y, Takehara Y, Kohriyama S, Sano Y, Tanabe K, Kobayashi T, Nakamura Y, Ohno T, Nozawa S 1991 Effects of prolactin on fertilization and cleavage of human oocytes. *Horm Res* 1:33-38
605. Messinis IE, Templeton AA 1987 Relationship between intrafollicular levels of prolactin and sex steroids and *in vitro* fertilization of human oocytes. *Hum Reprod* 2:607-609
606. Irahara M, Azuma K, Yamano S, Aono T 1991 Effect of prolactin in serum and follicular fluid on fertilization and cleavage of human oocyte. *Horm Res* 1:45-46
607. Randall G, Awadalla S, Shivers C 1990 Isolation, *in vitro* maturation, and fertilization of germinal vesicle oocytes obtained from the intact murine ovary. *J In Vitro Fert Embryo Transfer* 7:314-320
608. Yokkaichiya T, Fukaya T, Hoshiai H, Yajima A 1988 Improvement of mouse embryo development *in vitro* by prolactin. *Tohoku J Exp Med* 155:241-246
609. Yoshida K, Lin JM, Otsuka H, Takashima M, Okamura Y 1987 The effect of prolactin on the early embryogenesis of mice *in vitro*. *Sangyo Ika Daigaku Zasshi* 9:181-186
610. Fukuda A, Noda Y, Mori T, Mori C, Hashimoto H, Hoshino K 1988 Effects of prolactin on gametes and zygotes during *in vitro* fertilization in mice. *J In Vitro Fert Embryo Transfer* 5:25-30
611. Dodds WG, Fowler J, Peykoff A, Miller KF, Friedman CI, Kim MH 1990 The effect of prolactin on murine *in vitro* fertilization and embryo development. *Am J Obstet Gynecol* 162:1553-1559
612. Roblero L, Garavagno A 1979 Effect of oestradiol-17 β and progesterone on oviductal transport and early development of mouse embryos. *J Reprod Fertil* 57:91-95
613. McCormack JT, Greenwald GS 1974 Progesterone and oestradiol-17 β concentrations in the peripheral plasma during pregnancy in the mouse. *J Endocrinol* 62:101-107
614. Rider V, Moscoso H, Rizzo M 1987 Anti-progesterone monoclonal antibodies affect early cleavage and implantation in the mouse by mechanisms that are influenced by genotype. *J Reprod Fertil* 79:33-43
615. O'Neill C 1985 Partial characterization of the embryo-derived platelet-activating factor in mice. *J Reprod Fertil* 75:375-380
616. Orozco C, Clark IQ, Cock IE, Clarke FM 1994 A description of the basic system of components in pregnant mice sera responsible for "early pregnancy factor" activity. *J Reprod Fertil* 100:291-297
617. Cavanagh AC, Morton H 1994 The purification of early-pregnancy factor to homogeneity from human platelets and identification as chaperonin 10. *Eur J Biochem* 222:551-560
618. Sueoka K, Dharmarajan AM, Miyazaki T, Atlas SJ, Wallach EE 1988 Platelet activating factor-induced early pregnancy factor activity from the perfused rabbit ovary and oviduct. *Am J Obstet Gynecol* 159:1580-1584
619. Morton H, Cavanagh AC, Athanasas PS, Quinn KA, Rolfe BE 1992 Early pregnancy factor has immunosuppressive and growth factor properties. *Reprod Fertil Dev* 4:411-422
620. Cavanagh AC, Morton H, Rolfe BE, Gidley-Baird A 1982 Ovum factor: a first signal of pregnancy? *Am J Reprod Immunol* 2:97-101
621. Athanasas PS, Quinn KA, Wong TY, Rolfe BE, Cavanagh AC, Morton H 1989 Passive immunization of pregnant mice against early pregnancy factor causes loss of embryonic viability. *J Reprod Fertil* 87:495-502
622. Athanasas PS, Morton H, Dunglison GF, Kaye PL 1991 Antibodies to early pregnancy factor retard embryonic development in mice *in vitro*. *J Reprod Fertil* 92:443-451
623. Astwood E, Greep R 1938 A corpus luteum-stimulating substance in the rat placenta. *Proc Soc Exp Biol Med* 38:713-716
624. Galosy S, Talamantes F 1995 Luteotropic actions of placental lactogens at midpregnancy in the mouse. *Endocrinology* 136:3993-4003
625. O'Neill C, Quinn P 1981 Interaction of uterine flushings with mouse blastocysts *in vitro* as assessed by the incorporation of (³H)uridine of the embryo-derived platelet-activating factor in mice. *J Reprod Fertil* 62:257-262
- 625a. Lim H, Paria BC, Das SK, Dinchul JE, Langenbach R, Trzaskos JM, Dey SK 1997 Multiple female reproductive failures in cyclooxygenase 2-deficient mice. *Cell* 91:197-208
626. Shah GV, Sheth AR 1979 Is prolactin involved in sperm capacitation? *Med Hypotheses* 5:909-914
627. Mori C, Hashimoto H, Hoshino K, Fukuda A, Noda Y, Mori T 1988 Influences of prolactin upon spermatogenesis and spermatozoa during *in vitro* fertilization in mice. *J In Vitro Fert Embryo Transfer* 5:61-66
628. Bartke A 1980 Role of prolactin in reproduction in male mammals. *Fed Proc* 39:2577-2581
629. Korach KS 1994 Insights from the study of animals lacking functional estrogen receptor. *Science* 266:1524-1527
630. Liu X, Robinson GW, Wagner KU, Garrett L, Wynshaw-Boris A, Hennighausen L 1997 Stat5a is mandatory for adult mammary gland development and lactogenesis. *Genes Dev* 11:179-186
- 630a. Udy GB, Towers RP, Snell RG, Wilkins RJ, Park S-H, Ram PA,

- Waxman DJ, Davey HW 1997 Requirement of Stat5b for sexual dimorphism of body growth rates and liver gene expression. *Proc Natl Acad Sci USA* 94:7239-7244
631. Horseman ND, Zhao W, Montecino-Rodriguez E, Tanaka M, Nakashima K, Engle SJ, Smith F, Markoff E, Dorskind K 1997 Defective mammapoiesis, but normal hematopoiesis, in mice with a targeted disruption of the prolactin gene. *EMBO J* 16: 6926-6935
632. Toscani A, Mettus RV, Coupland R, Simpkins H, Litvin J, Orth J, Hatton KS, Reddy EP 1997 Arrest of spermatogenesis and defective breast development in mice lacking *A-myb*. *Nature* 386: 713-717
633. Fantl V, Stamp G, Andrews A, Rosewell I, Dickson C 1995 Mice lacking cyclin D1 are small and show defects in eye and mammary gland development. *Genes Dev* 9:2364-2372
634. Freemark M, Nagano M, Ederly M, Kelly PA 1995 Prolactin receptor gene expression in the fetal rat. *J Endocrinol* 144:285-292
635. Brown-Borg HM, Zhang FP, Huhtaniemi I, Bartke A 1996 Developmental aspects of prolactin receptor gene expression in fetal and neonatal mice. *Eur J Endocrinol* 134:751-757